

## Understanding the causes and consequences of past marine carbon cycling variability through models



Dominik Hülse<sup>a,\*</sup>, Sandra Arndt<sup>a,d</sup>, Jamie D. Wilson<sup>a</sup>, Guy Munhoven<sup>b</sup>, Andy Ridgwell<sup>a,c</sup>

<sup>a</sup> BRIDGE, School of Geographical Sciences, University of Bristol, Bristol BS8 1SS, UK

<sup>b</sup> Institute of Astrophysics and Geophysics, University of Liège, Liège B-4000, Belgium

<sup>c</sup> Department of Earth Sciences, University of California, Riverside, CA 92521, USA

<sup>d</sup> Department Geosciences, Environment and Society, Université Libre de Bruxelles, Brussels, Belgium

### ARTICLE INFO

#### Keywords:

Biological carbon pump  
Earth system models  
Ocean biogeochemistry  
Marine sediments  
Paleoceanography

### ABSTRACT

On geological time-scales, the production and degree of recycling of biogenic carbon in the marine realm and ultimately its removal to sediments, exerts a dominant control on atmospheric CO<sub>2</sub> and hence variability in climate. This is a highly complex system involving a myriad of inter-connected biological, chemical, and physical processes. For this reason alone, linking observations, often highly abstracted in the form of proxies, to the primary processes involved and ultimately to explanatory hypotheses for specific geological events and transitions, is challenging. The past few decades have seen a progressive improvement in theoretical and process-based understanding of the various components that make up the marine carbon cycle and, hand-in-hand with this, the development of numerical model representations of the complete system. Models have also been designed and/or adapted with paleoclimate questions in mind and applied to quantitatively explore the role of the marine carbon cycle in both perturbations and long-term geologic evolutionary trends in global climate, and possible feedbacks between them. However, we must ask whether paleoclimate models incorporate sufficiently appropriate representations of the dynamics and sensitivities of the marine carbon cycle, and indeed, whether in the geological context, we really know what these dynamics are.

Here we provide a comprehensive overview of how marine carbon cycling and the biological carbon pump is treated in available paleoclimate models, with the aim of critically evaluating their ability to help interpret past marine carbon cycle and climate dynamics. To this end, we first provide an overview of commonly used paleoclimate models and some of their associated paleo-applications, drawing from a wide range of global carbon cycle box models and Earth system Models of Intermediate Complexity (EMICs). Secondly, we review and evaluate the three dominant processes involved in the cycling of organic and inorganic carbon in the marine system and how they are represented in models, namely: biological productivity at the ocean surface, remineralisation/dissolution of particulate carbon within the water column, and the benthic-pelagic coupling at the seafloor. We generate and employ illustrative examples using the model GENIE to show how different parameterisations of water-column and sediment processes can lead to significantly different model projections. Our compilation reveals the prevalence of static parameterisations of marine carbon cycling among existing paleoclimate models, which are commonly empirically derived from present-day observations. Although such approaches tend to represent carbon transfer in the modern ocean well, they are potentially compromised in their ability to reflect the true degree of freedom and strength of feedbacks with respect to past climate events, particularly those characterised by environmental boundary conditions that differ fundamentally from today. Finally, we discuss the importance of using models of different complexities and how questions of model uncertainty may start to be addressed.

### 1. Introduction

On a wide range of geologically-relevant and resolvable time-scales, from 10<sup>3</sup> to > 10<sup>6</sup> years, the climate system is thought to be largely

driven by variations in atmospheric CO<sub>2</sub>, itself controlled by a dynamic balance between varying CO<sub>2</sub> inputs from volcanoes and metamorphic alteration of rock, and the removal of CO<sub>2</sub> from the ocean-atmosphere system through weathering and burial of carbon in marine sediments

\* Corresponding author.

E-mail address: [Dominik.Huelse@bristol.ac.uk](mailto:Dominik.Huelse@bristol.ac.uk) (D. Hülse).




Fig. 1. A simplified view of the global carbon cycle showing the approximate carbon stocks in Pg C and main annual fluxes in Pg C year<sup>-1</sup> for the preindustrial era (largely based on IPCC 2013 (Ciais et al., 2014) and Hain et al. (2014b), GPP: gross primary production). Years in red indicate the time-scales at which the reservoirs exert control on atmospheric CO<sub>2</sub> concentrations. Thus, the biological carbon pump in the ocean is important on time-scales shorter than about 1000 years, whereas the sediments become influential on time-scales of 1000 to 100,000 years.

(Fig. 1; Berner, 1991; Berner and Caldeira, 1997; Kump et al., 2000; Ridgwell and Zeebe, 2005). A key motivator of paleoclimate research is hence to understand the controls, characteristic time-scales and the dynamic balance between carbon sources and sinks as well as its possible feedbacks with global climate. This endeavour rests heavily on numerical models to help quantify potential dynamic behaviours and sensitivities of responses and to test hypotheses. To be successful however, these models must include appropriate representations of the key carbon cycle processes and/or exhibit real-world relevant dynamics. To what degree this is true is the topic of this review which we focus around marine carbon cycling and in particular the sequence of processes that create and transfer carbon from the ocean surface to the underlying sediments.


In the modern carbon cycle, the net transfer of CO<sub>2</sub> from the atmosphere to the oceans and then sediments, is primarily a direct consequence of the combined effect of the solubility and biological pump (Volk and Hoffert, 1985). The solubility pump describes the exchange of CO<sub>2</sub> between ocean surface and atmosphere, and its redistribution as dissolved inorganic carbon within the ocean interior by large-scale circulation. It is estimated to account for only ~10% of the surface-to-deep gradient of (preindustrial) dissolved inorganic carbon (DIC) (Sarmiento and Gruber, 2006) and cannot itself lead to carbon removal from the surficial system. As its potential for modifying atmospheric CO<sub>2</sub> is limited and it mostly acts on a sub-kyr time-scale, we do not address the paleo role of this process in any detail in this paper. The biological carbon pump, for which we will expand on more common definitions for our geological context, refers here to both the organic and inorganic carbon fixed by primary producers in the euphotic zone, the efficiency of its transfer to the underlying ocean interior, and the efficiency with which it is buried in the underlying sediments (Fig. 2; Volk and Hoffert, 1985). The growth of phytoplankton in the light-flooded surface layer of the global ocean removes nutrients and carbon from the water and transforms them into cellular material and/or inorganic carbonates. Upon organism death, part of this produced particulate organic and inorganic carbon (POC, PIC) sinks out of the euphotic zone. In addition, dissolved organic carbon (DOC) is either scavenged by particle aggregates and sinks to the deep ocean or is redistributed into the deeper layers by ocean circulation. In today's well oxidised ocean a large fraction of the gross primary production of organic matter is remineralised by bacterial activity in the water column and < ~0.5% of it is ultimately buried in marine sediments (Fig. 1;

Hedges and Keil, 1995; Burdige, 2007).

Despite intense ocean upwelling and mixing, the biological pump results in distinct ocean depth gradients of carbon, nutrients and oxygen. In particular its control on surface ocean DIC concentration, which is next to temperature, wind speed and ambient pH one of the main drivers of air-sea carbon dioxide fluxes, is of fundamental importance for the global carbon cycle and thus global climate (Volk and Hoffert, 1985). In fact, modelling studies have shown that atmospheric CO<sub>2</sub> concentrations would be significantly higher above an abiotic ocean (Sarmiento and Toggweiler, 1984; Archer et al., 2000b). Sarmiento and Gruber (2006) show that the organic part of the biological pump is responsible for about 70% and the inorganic part for 20% of the total preindustrial dissolved inorganic carbon transfer from the surface to the deeper ocean. (The remaining 10% are attributed to DIC variations driven by temperature variations, ocean mixing and the solubility pump.)

The efficiency of the biological pump depends not only on the rate of carbon fixation and export out of the surface layer, but also on the depth at which the organic and inorganic carbon is respired or dissolved. This depth determines the time during which carbon is isolated from the atmosphere and therefore atmospheric CO<sub>2</sub> concentration exhibits an inverse relationship to the efficiency of the biological pump (Yamanaka and Tajika, 1996; Boyd and Trull, 2007). Estimates for global organic carbon exported from the surface to the deeper ocean are in the range of 5–11 Pg C year<sup>-1</sup> (Laws et al., 2000; Dunne et al., 2007; Henson et al., 2011). In the contemporary, well-oxygenated ocean, only a small fraction of the carbon organically produced in the surface ocean escapes microbial degradation (remineralsation) and is eventually buried in the sediment (e.g. Middelburg and Meysman, 2007). The inorganic carbon flux is also strongly coupled to the organic carbon flux by the biologically driven carbonate precipitation and the effect of organic matter remineralisation on carbonate preservation (see Box 1 or, e.g., Hales, 2003; Ridgwell and Zeebe, 2005).

The complex interplay and strong feedback between fluxes and transformations of carbon in the ocean-sediment system limit our predictive ability of the global climate system and its evolution throughout Earth's history and critically disguise the underlying physical and biogeochemical process interplay behind our paleo observations that simply reflect the net process outcome (Falkowski et al., 2000). Appropriate mathematical models can help as they allow us to elucidate the various rates of biogeochemical processes and related fluxes that



**Fig. 2.** Schematic of the main processes constituting the ocean's solubility and biological pumps (labelled 1 to 6). 1: Air-sea gas exchange of  $\text{CO}_2$ , used by phytoplankton to form cellular material. Through complex recycling and aggregation pathways, agglomerations of POC and PIC are formed, heavy enough to sink out of the euphotic zone. 2: Precipitation of  $\text{CaCO}_3$  by, e.g., coccolithophores and foraminifera in the open ocean or by corals and coralline algae in the shallow-water contributing to the formation of reefal structures, resulting in higher  $p\text{CO}_2$  at the surface and therefore a transfer of  $\text{CO}_2$  to the atmosphere ( $\text{Ca}^{2+} + 2\text{HCO}_3^- \rightarrow \text{CaCO}_3 + \text{CO}_2(\text{aq}) + \text{H}_2\text{O}$ ). 3: Carbonate sinking into the deeper ocean and forming deep-sea sediments. At the lysocline the rate of dissolution increases dramatically until below the CCD no  $\text{CaCO}_3$  is present in the sediment any more. 4: Vertical settling flux of POC (in combination with PIC) that decreases approximately exponentially with depth. 5: Deposition of organic carbon in the deep sea. Only a small fraction escapes degradation and is buried in the sediments. 6: DIC in the upper ocean is created through degradation and egestion processes and upwelling of DIC rich subsurface waters. These processes contribute to raise surface water  $p\text{CO}_2$ , driving a transfer of  $\text{CO}_2$  to the atmosphere.

### Box 1

#### Calcium carbonate thermodynamics primer.

One of the most common minerals on Earth is calcium carbonate ( $\text{CaCO}_3$ ) which is almost exclusively formed by biological processes. In the ocean  $\text{CaCO}_3$  most commonly exists in two crystal forms: *calcite* and *aragonite*. Because calcite is the more thermodynamically stable phase it is more abundant in the ocean and forms almost all deep sea carbonate sediments. Aragonite, on the other hand, is found in shallow-water sediments (e.g. corals). Carbonate precipitation can be described by the following chemical reaction:  $\text{Ca}^{2+} + 2\text{HCO}_3^- \rightarrow \text{CaCO}_3 + \text{CO}_2(\text{aq}) + \text{H}_2\text{O}$

Whether  $\text{CaCO}_3$  precipitates or dissolves can be directly linked to the concentrations of  $\text{Ca}^{2+}$  and  $\text{CO}_3^{2-}$  in sea-water which controls the stability of its crystal structure (Ridgwell and Zeebe, 2005). This is expressed as the *saturation state*  $\Omega$  of the solution and is defined as

$$\Omega = [\text{Ca}^{2+}] \times [\text{CO}_3^{2-}] / K_{\text{sp}}$$

where  $K_{\text{sp}}$  is a solubility constant which scales with increasing pressure (Zeebe and Wolf-Gladrow, 2001). Calcium carbonate is thermodynamically favourable to precipitate at  $\Omega > 1$  (i.e. in oversaturated environments) and prone to dissolution when  $\Omega$  is smaller than unity. The saturation state of the ocean generally decreases with water depth in response to the combined effects of the pressure-induced increase in the solubility constant and the release of  $\text{CO}_2$  through the degradation of sinking POC that decreases the ambient pH and thus the carbonate ion concentration (Ridgwell and Zeebe, 2005; Soetaert et al., 2007). Oceanic waters become undersaturated (i.e.  $\Omega < 1$ ) below the depth of the *saturation horizon* (i.e.  $\Omega = 1$ ) and carbonates start to dissolve. However, the process of dissolution proceeds only extremely slowly in the beginning. The depth at which the dissolution rate increases considerably and impacts become noticeable is known as the *lysocline* (Broecker, 2003) (sometimes defined at  $\Omega = 0.8$ , Milliman et al., 1999). Deeper still, the rate of dissolution becomes fast enough to exactly balance the rate of supply of carbonates to the sediments and therefore no carbonates are preserved. This is termed the *carbonate compensation depth* (CCD, compare Fig. 2).

comprise the net result (ideally, but rarely, quite equal to the observation). They also present the opportunity to formally and quantitatively test hypotheses that arise from observations. Furthermore, they allow us to explore and discover behaviours and dynamics that may not necessarily have ever occurred in Earth history, or more usefully, behaviours and dynamics that we did not a priori suspect were taking place until their possibility was revealed by a model.

The great challenge in successfully applying paleoclimate models arises because of practical temporal and spatial constraints on computation, unconstrained and uncertain boundary conditions, but also simply because of the lack of mechanistic knowledge. It is impossible therefore for paleoclimate (or indeed, any) models to capture the full complexity of the complete carbon cycle, at least not perfectly (Cox et al., 2000; Friedlingstein et al., 2003, 2006). Simplification is necessary and ideally would be based on a sound mechanistic understanding of the underlying processes and be consistent and traceable between models. Instead, the choices of how to parameterise processes and which processes to neglect altogether, vary between models and can depend on the specific questions or paleo events being addressed. It is hence critical that we have a good understanding of the consequences of these choices and what the implications are for the degree of confidence we can draw from model results when applied in the paleo realm. Particular care is needed in respect of the applicability of parameterisations that are derived from observations of the present-day ocean-sediment system because environmental conditions may be very different during past extreme events (compare Fig. 3).


Unlike in the case of modern (IPCC, Randall et al., 2007) and past climate dynamics (PMIP, whose aim is to test the ability of General Circulation Models (GCMs) to reproduce feedbacks and large changes in the climate system, e.g. Kageyama et al., 2016; Lunt et al., 2017), there has not yet been any systematic review study of the marine carbon cycle dynamics represented in paleoclimate models, a process which could help to substantially upgrade the performance of these models and raise the degree of confidence in their results. The geologic context also requires sediment dynamics to be considered, a component not typically considered as part of the biological carbon pump, as benthic-pelagic coupling plays an important role for carbon cycling and Earth's recovery from global warming events, especially on longer time-scales (> 1000 years; Fig. 1). Conversely, to simplify the analysis, we omit shorter time-scale processes that impact upon the biological carbon pump such as ocean circulation, which at least in the modern/future context, are discussed in detail elsewhere (e.g. IPCC, Ciais et al., 2014). Furthermore, this study does not aim to provide an exhaustive review of all critical aspects of ocean biogeochemistry, particularly since facets such as the marine iron cycle and its influence on the biological pump are relatively poorly studied and represented even in models of the

modern ocean (Tagliabue et al., 2016). Instead, we focus on the core behaviours and dynamics of three key aspects of marine organic and inorganic carbon cycling and their encapsulation into Earth system models: biological productivity at the ocean surface, remineralisation/dissolution of carbon within the water column and the benthic-pelagic coupling at the seafloor. The crux of the issue addressed here is then the range of formulations used to represent these key carbon cycle processes and the resulting divergence of predictions for carbon cycle change under different boundary conditions, such as increased global temperatures, ocean anoxia and euxinia (Fig. 3).

The structure of this review is as follows. First, after a brief summary of the history of model development (as it helps to understand where many of the widely used carbon cycle parameterisations originally come from and how and why they were developed), different types of paleoclimate models are presented and compared; and an overview for their characteristic time-scales of application is given (Section 2). Thereafter, the focus of the paper is on reviewing the three key processes of the biological carbon pump and evaluating their representation in paleoclimate models by testing their ability to predict the response of carbon cycling under different extreme climate conditions (Section 3). The final part (Section 4), highlights the importance for applying models of different complexities, and summarises two key outstanding modelling issues.

## 2. Overview of numerical models commonly employed in paleoclimate research

Numerical models are essential tools that help us to understand the complex Earth system dynamics, including how climate and the biological pump have changed in the past and how they may evolve in the future (see e.g. Alverson et al., 2003; McGuffie and Henderson-Sellers, 2005; Randall et al., 2007). Existing Earth system models vary considerably in complexity from highly parameterised conceptual models (e.g., box models) to comprehensive coupled General Circulation Models (GCMs), with different scientific questions often being tackled with different model approaches. Writing a fully comprehensive review of all available models including a representation of the biological pump would be an impossible task and quickly out of date. We have hence chosen to focus on more computationally efficient models, such as box models (Section 2.2) and intermediate complexity models (Section 2.3), which include a minimum set of components representing the biological carbon pump. Furthermore, the particular models we include have all been used to study biogeochemical cycling for some interval in Earth history and have generated simulations that span order  $10^4$  years or more (and thus are able to model processes involving the deep ocean and the marine sediments). We hence omit higher



**Fig. 3.** Schematic representation of the potential functioning of the biological pump in the modern world (left), during the Eocene (middle) and the oceanic anoxic event 2 (right, OAE2). There is much evidence that global mean ocean temperatures exceeded those of the modern day by several degrees during the Eocene epoch (55.5–33.7 Ma) and theory predicts that elevated temperatures should cause a decrease in the efficiency of the biological pump as sinking POC is more quickly remineralised. In contrast, during OAE2 (ca. 93.5 Ma) the ocean exhibited widespread oxygen depletion and photic zone euxinia (occurrence of hydrogen sulphide ( $H_2S$ )). At the same time the marine sediments are characterised by enhanced deposition of organic carbon-rich black layers.

**Table 1**  
Overview of EMICs mentioned in this study and their characteristics related to the biological pump.

Models	Ocean Model	Ocean Carbon Cycle			Sediments
	Resolution	Surface Production	Tracers	Degradation	
<b>EMICs</b>					
<b>Bern 2.5D</b>	Zonally averaged, 3-basin circulation model, 14 vertical layers Marchal et al. (1998)	Michaelis-Menten kinetics, PO <sub>4</sub> limiting nutrient and relation to temperature. Marchal et al. (1999)	DOC, DIC, C, δ <sup>13</sup> C, δ <sup>14</sup> C, PO <sub>4</sub> , ALK, O <sub>2</sub> , CaCO <sub>3</sub> ; Marchal et al. (1998b)	Power law (Martin)	No sediment representation
<b>Bern 3D</b>	3-D, horizontal 36x36 equal area boxes, 10°x(3.2-19.2)°, 32 vertical layers; Müller et al. (2006)	Fully coupled carbon cycle, Michaelis-Menten nutrient uptake kinetics, limited by PO <sub>4</sub> , Fe and light. Parekh et al. (2008); Tschumi et al. (2008)	CFC-11, PO <sub>4</sub> , DOP, DIC, DOC, δ <sup>13</sup> C, δ <sup>14</sup> C, δ <sup>39</sup> Ar, Ar, ALK, O <sub>2</sub> , FeT; Müller et al. (2008), Parekh et al. (2008)	Power law (Martin)	Vertically integrated, dynamic model of top 10cm: oxic respiration and denitrification. CaCO <sub>3</sub> , opal, POM, clay and DIC, ALK, PO <sub>4</sub> , NO <sub>3</sub> , O <sub>2</sub> , silicic acid; Tschumi et al. (2011)
<b>CLIMBER-2</b>	3 zonally averaged basins, 2.5° (meridional), 20 vertical layers	Fully coupled carbon cycle (HAMOCC3) + ecosystem model, Six and Maier-Reimer (1996)	DIC, DOC, PO <sub>4</sub> , O <sub>2</sub> , Alkalinity, silicate, isotopes ( <sup>12</sup> C, <sup>13</sup> C, <sup>14</sup> C, <sup>39</sup> A), δ <sup>18</sup> O; Brovkin et al. (2002)	Power law (Suess)	CaCO <sub>3</sub> model with oxic-only sediment respiration (Archer, 1996a)
<b>GENIE</b>	3-D, horizontal 36x36 equal area boxes, 10° x (3.2-19.2)°, 8/16 vertical layers	Fully coupled carbon cycle, Michaelis-Menten nutrient uptake kinetics, limited by PO <sub>4</sub> , NO <sub>3</sub> , Fe and light, Ridgwell et al. (2007)	Ca. 56 dissolved and various solid tracers; see www.seao2.info/mycgenie.html	Double Exponential / Temp dependent / Power Law / Ballasting	Vertically integrated model: u.a. CaCO <sub>3</sub> , detrital material, δ <sup>13</sup> C, δ <sup>14</sup> C; Ridgwell and Hargreaves (2007)
<b>GEOCLIM</b>	3 zonally averaged basins, 2.5° (meridional), 20 vertical layers (uses CLIMBER-2)	Fully coupled carbon cycle model, PO <sub>4</sub> limiting nutrient (using COMBINE, Goddérís and Joachimski (2004))	PIC, POC, PO <sub>4</sub> , O <sub>2</sub> , DIC, ALK, pH, δ <sup>13</sup> C, δ <sup>18</sup> O, Ca <sup>2+</sup> , CaCO <sub>3</sub>	Recycling rate linear function of [O <sub>2</sub> ]	Simple org. C, P model (COMBINE): oxic layer (C:P ratio ~ [O <sub>2</sub> ]) and constant sulfate reduction zone; Goddérís and Joachimski (2004)
<b>LOVECLIM</b>	3-D, 3° x 3°, 30 vertical layers; Goosse et al. (2010)	Fully coupled carbon cycle; LOCH:Mouchet and Francois (1996); rain ratio depends on silica, tmp, CaCO <sub>3</sub> )	DIC, Alkalinity, dissolved inorganic phosphorous, DOC, POC, O <sub>2</sub> , org. and inorg. δ <sup>13</sup> C, δ <sup>18</sup> O, CaCO <sub>3</sub> , silica, opal	Power law (Martin)	Constant part of POC & PIC is preserved; Goosse et al. (2010)
<b>MESMO</b>	3-D, horizontal 36x36 equal area boxes, 10° x (3.2-19.2)°, 16 vertical layers	Fully coupled carbon cycle, Michaelis-Menten nutrient uptake kinetics, limited by PO <sub>4</sub> , NO <sub>3</sub> , CO <sub>2</sub> , Fe, light and tmp; Matsumoto et al. (2008)	CFC-11, NO <sub>3</sub> , N*, <sup>15</sup> N, PO <sub>4</sub> , O <sub>2</sub> , DIC, CaCO <sub>3</sub> , ALK, POC, DOC, δ <sup>13</sup> C, <sup>14</sup> C	Temp dependent	Vertically integrated model: u.a. CaCO <sub>3</sub> , detrital material, δ <sup>13</sup> C, δ <sup>14</sup> C; Ridgwell and Hargreaves (2007)
<b>MoBidIC</b>	3 zonally averaged basins, 5° (meridional), 19 vertical layers	Fully coupled carbon-cycle, PO <sub>4</sub> limiting nutrient, Michaelis-Menten kinetics, Crucifix, (2005)	DIC, DI <sup>13</sup> C, DOC, DO <sup>13</sup> C, ALK, PO <sub>4</sub> , O <sub>2</sub> , <sup>14</sup> C, POC, CaCO <sub>3</sub>	Single exponential	No sediment representation
<b>UVic</b>	3-D, 1.86° (meridional) x 3.6° (zonal), 19 vertical layers	Fully coupled carbon cycle + NPZD ecosystem representation + dynamic iron cycle; e.g. Schmittner et al. (2008), Keller et al. (2012), Nickelsen et al. (2015)	Phytoplankton (nitrogen fixers and others), NO <sub>3</sub> , PO <sub>4</sub> , O <sub>2</sub> , DIC, CaCO <sub>3</sub> , ALK, zooplankton, particulate detritus, POC, DOC, δ <sup>13</sup> C, δ <sup>18</sup> O, Fe, Fe <sub>p</sub>	Single exponential	CaCO <sub>3</sub> model with oxic-only sediment respiration (Archer, 1996a)

<b>Glossary</b>	Circulation & Temp feedback; reduced complexity atm. GCM	Ecological model	Double exponential / Temp dependent	Vertically integrated, organic and inorganic carbon burial
	Circulation & Temp feedback; highly simplified or basic dynamics atm.	Export production model (10 or more tracers)	Power law or single exponential	Vertically integrated inorganic carbon burial - no organic carbon burial
	No or simplified feedbacks (e.g. weathering)	Export production model (less than 10 tracers) or simpler model	Linear/fixed fraction	Constant preservation

resolution ocean models and fully coupled GCMs, that while encompassing sophisticated representations of ocean biogeochemistry and circulation and describing many details of fluxes between the ocean-atmosphere systems (see e.g. Washington and Parkinson, 2005), generally focus on time scales of a few thousands of years or less, and generally do not take into account sediment processes (examples of GCMs that have been used for paleo studies with a focus on the biological pump are: MPIOM, Heinze and Ilyina (2015) and CCSM3,

Winguth et al. (2012)).

Historically, paleoclimate models with a focus on the ocean's biological pump have evolved along two distinct paths. Global carbon cycle box models were designed around simplified multi-box or advection-diffusion models of the ocean circulation and global carbon cycling, while Earth system Models of Intermediate Complexity (EMICs; Claussen et al., 2002) can be considered as coarse resolution Earth system models (e.g. Eby et al., 2013). We hence make a distinction

Table 2

Overview of global carbon cycle box models mentioned in this study and their characteristics related to the biological pump (glossary of Table 1 applies here as well).

Models	Ocean Model	Ocean Carbon Cycle			Sediments
	Resolution	Surface Production	Tracers	Degradation	
<b>Box models</b>					
<b>BICYCLE</b>	10 homogeneous boxes (5 surface, 2 intermediate, 3, deep); Köhler et al. (2005)	Fixed fraction of PO <sub>4</sub> , Munhoven and François (1996)	PIC, POC, PO <sub>4</sub> , O <sub>2</sub> , DIC, ALK, δ <sup>13</sup> C, δ <sup>14</sup> C	Fixed fraction	No sediment representation or restoration of prescribed lysocline or [CO <sub>3</sub> <sup>2-</sup> ]
<b>CYCLOPS</b>	14/13 homogeneous boxes, e.g. Keir 1988, 1990; later 18 boxes, Hain et al. 2010	Fixed fraction of PO <sub>4</sub> depending on latitude, Keir (1988)	PO <sub>4</sub> , O <sub>2</sub> , DIC, ALK, δ <sup>13</sup> C, δ <sup>14</sup> C, POC, CaCO <sub>3</sub>	Fixed fraction depending on latitude	1. Similar to LOSCAR (but const. porosity); 2. Respiration driven CaCO <sub>3</sub> dissolution, Sigman et al. (1996)
<b>DCESS</b>	2 x 55 boxes, 1 hemisphere, 100m vertical resolution, Shaffer et al. 2008	Export production model, PO <sub>4</sub> limiting nutrient, Yamanaka and Tajika (1996)	POC, PO <sub>4</sub> , O <sub>2</sub> , DIC, ALK, δ <sup>12,13,14</sup> C, CaCO <sub>3</sub>	Single exponential	Semi-analytical, oxic respiration and denitrification, CaCO <sub>3</sub> , Shaffer et al. 2008
<b>LOSCAR</b>	10 homogeneous boxes for modern ocean (13 e.g. for the P/E-version)	Biological uptake using Michaelis-Menten kinetics, PO <sub>4</sub> limiting nutrient	TCO <sub>2</sub> , TA, PO <sub>4</sub> , O <sub>2</sub> , δ <sup>13</sup> C, pH, Mg/Ca	Fixed fraction	%CaCO <sub>3</sub> in the bioturbated layer as a fct. of sediment rain, dissolution, burial and chemical erosion - also variable porosity
<b>MBM</b>	10 homogeneous boxes (5 surface, 2 intermediate, 3, deep)	Similar to BICYCLE	DIC, TA, PO <sub>4</sub> , O <sub>2</sub> , CO <sub>2</sub> , δ <sup>13</sup> C, δ <sup>14</sup> C, OM, calcite, aragonite	Power law (Martin)	Advection-diffusion-reaction model MEDUSA: solids (clay, calcite, aragonite, OM), solutes (CO <sub>2</sub> , HCO <sub>3</sub> <sup>-</sup> , CO <sub>3</sub> <sup>2-</sup> , O <sub>2</sub> )
<b>PANDORA</b>	10/11 homogeneous boxes, e.g. Broecker and Peng (1986, 1987)	Limiting nutrients (PO <sub>4</sub> , NO <sub>3</sub> ) with diff. residence times per box	PO <sub>4</sub> , NO <sub>3</sub> , O <sub>2</sub> , DIC, ALK, δ <sup>13</sup> C, δ <sup>14</sup> C, POC, silica, CaCO <sub>3</sub> , <sup>39</sup> Ar, <sup>3</sup> He, <sup>13</sup> C/ <sup>12</sup> C	Fixed fraction	No sediment representation
<b>SUE</b>	Uses set-up of Bern 2.5D or PANDORA model	Export production model (siliceous and non-siliceous phytoplankton), limiting nutrients (PO <sub>4</sub> , H <sub>4</sub> SiO <sub>4</sub> , Fe); Ridgwell (2001)	PO <sub>4</sub> , H <sub>4</sub> SiO <sub>4</sub> , Fe, DIC, ALK, O <sub>2</sub> , CaCO <sub>3</sub> , POC, opal, DOM, δ <sup>12,13</sup> C, δ <sup>16,17,18</sup> O	Single exponential	CaCO <sub>3</sub> and opal as fct. of saturation state, Ridgwell et al. (2002), Archer (1991)

between the two and discuss them largely separately. Box models and EMICs can also mostly be separated by the general absence of explicit climate feedback with carbon cycling in the former, and inclusion in the latter (see Tables 1 and 2), although most EMICs include few of the atmospheric feedbacks (e.g. internal climate oscillations such as ENSO, changes with climate in wind strength and zonal position, cloud cover and albedo) that fully coupled GCMs can be thought of as specializing in.

### 2.1. Model development history

The model-supported analysis of global carbon cycling started in the mid-fifties with the box-models of Eriksson and Welander (1956), Craig (1957) and Bolin and Eriksson (1959). Welander (1959) further added a diffusive representation of carbon transport in the sea, thus producing the first so-called box-diffusion models. Garrels and MacKenzie (1972) adopted the box-modelling framework to the steady-state long-term rock cycle. Time-dependent quantitative simulation experiments about the role of the carbonate and silicate rock cycles for the evolution of atmospheric CO<sub>2</sub> were realized with the BLAG model (Bernier et al., 1983). This was the first in a long line of box-models often referred to as *Berner models* (Bernier, 1991, 1994; Bernier and Kothavala, 2001).

Box-diffusion models and variants thereof became widely used in the seventies and early eighties to study the partitioning of anthropogenic CO<sub>2</sub> between the atmosphere and the sea (Oeschger et al., 1975; Broecker et al., 1980; Hoffert et al., 1981; Siegenthaler, 1983). Early investigations on the glacial-interglacial CO<sub>2</sub> variations documented in the ice-cores relied again on simple three- to four-box models (Berger, 1982; Keir, 1983; Keir and Berger, 1983; Sarmiento and

Toggweiler, 1984; Siegenthaler and Wenk, 1984; Knox and McElroy, 1984, to cite only a few pre-1985 papers). These models were also the first ones to explicitly consider the surface to deep-sea transport of carbon by biogenic particles. The term “biological pump” was also coined at about that time by Volk and Hoffert (1985). Later, configurations with a larger number of boxes came into usage: the 13-box CYCLOPS (Keir, 1988); the ten-box PANDORA (Broecker et al., 1986); the ten-box model of Munhoven and François (1994, 1996); and the 19-box model of Michel et al. (1995) (only oceanic reservoirs considered). Including a larger number of boxes helped on the one hand to resolve the major ocean basins and water masses, but, on the other hand, increased the number of free parameters in the models (mainly the water exchange fluxes between the reservoirs) and thus made them more and more difficult to calibrate. In the mean time, the first three-dimensional ocean carbon cycle model had made its appearance (Maier-Reimer and Hasselmann, 1987; Bacastow and Maier-Reimer, 1990). The physically-based water circulation field that these models were based upon could be used to overcome the calibration dilemma of the multi-box models mentioned above (Munhoven and François, 1994, 1996; Michel et al., 1995). Box-models fell somewhat in disgrace after Archer et al. (2000b) had shown that they had a considerably higher sensitivity to high-latitude changes than two- or three-dimensional models, questioning the applicability of box-models for the quantitative assessment of the effects of high-latitude changes of biogeochemical and physical processes on atmospheric p CO<sub>2</sub>. Lane et al. (2006) finally demonstrated that the high-latitude sensitivity diagnosed by Archer et al. (2000b) is not related to the model type (box, versus 2D or 3D), but rather to the way the models are constructed. During the nineties, faster processors and improved data storage devices progressively allowed the application of

increasingly resolved models with increasingly improved physics on geological relevant time scales. In particular, coupled, multi-dimensional paleoclimate models, so-called Earth system Models of Intermediate Complexity (EMICs) have emerged in response to the rapid increase in computer power. As a result, box-models have to some extent been superseded by EMICs, especially, for the study of scientific problems that involve changing water mass distributions, which are difficult to take into account in a consistent and mechanistic way in box-models, or if vertical re-distributions of fluxes or solute materials must be tracked. Box-models nevertheless remain indispensable tools to analyse and interpret the results of more complex 2D and 3D models (Meinshausen et al., 2011) and to improve our general process understanding (e.g. Marinov et al., 2008). They also continue to be used as research tools on their own for time scales of hundreds of millions of years (e.g., Arvidson et al., 2014), hundreds of thousands of years (e.g., Hain et al., 2014a; Frieling et al., 2016) and future evolution of global carbon cycle (e.g., Boudreau et al., 2010; Köhler, 2016). Finally, they also continue to be developed (e.g., Zeebe, 2012). The Garrels and MacKenzie (1972) model was even revised and extended after more than thirty years to yield a new model, called MAGic (Arvidson et al., 2006). MAGic is unique in that it uses thermodynamic constants for its pH calculations that are based upon Pitzer equations, which are not biased towards the present-day sea salt composition, but can cope on varying composition.

## 2.2. Global carbon cycle box models

Global carbon cycle box models (GCCBMs) are a very conceptualised representations of the Earth system that focus on the global carbon cycle and are employed to test hypotheses and to quantify large scale processes. Due to their high computational efficiency they allow the investigation of long-term dynamics in the Earth system and its biogeochemical evolution (compare Fig. 4). Sarmiento and Toggweiler (1984) for instance, simulated with a simple 4-box model of the ocean-atmosphere the substantial influence of high latitude surface ocean productivity and thermohaline overturning rate on atmospheric CO<sub>2</sub> fluctuations during glacial-interglacial cycles. However, GCCBMs incorporate highly simplified representations of both the atmosphere and ocean systems with the notable exception of the GEOCLIM family of models which use the biogeochemical ocean-atmosphere model COMBINE (Goddéris and Joachimski, 2004) and couple it to the three-dimensional climate model FOAM (Donnadieu et al., 2006). In general, the atmosphere is encapsulated into a simple one-box model and atmospheric processes are reduced to simple gas and sometimes heat exchange (e.g., DCESS Shaffer et al., 2008). Without a fully-resolved atmosphere, the upper oceanic boundary conditions (wind stress, heat flux, equilibrium state with fixed boundary conditions and fresh-water flux) can also not be defined in a physically rigorous way. Therefore,

the representation of ocean circulation in these models is highly parameterised and ranges from simple multi-box models with prescribed exchange fluxes (e.g., BICYCLE, Köhler et al., 2005) to advection-diffusion box models (e.g. DCESS, Shaffer et al., 2008; GEOCLIM reloaded, Arndt et al., 2011). Ice sheet dynamics are generally neglected due to the simplicity of the ocean-atmosphere models. Terrestrial vegetation and weathering fluxes are generally integrated in form of simplified rate laws (e.g., BICYCLE; Köhler et al., 2005).

## 2.3. Earth system Models of Intermediate Complexity

Earth system Models of Intermediate Complexity (EMICs) have been developed to close the gap between the computationally efficient, but conceptual box models and the computationally expensive General Circulation Models (Claussen et al., 2002). The first EMIC-like models that already included and coupled important parts of the Earth system were developed in the early 1980s (Petoukhov, 1980; Chalikov and Verbitsky, 1984). The spatial resolution of EMICs is coarser than that of “state-of-the-art” GCMs. However, they explicitly simulate, in an interactive mode, the basic components of the Earth system, including the atmosphere, ocean, cryosphere and land masses, over a very wide range of temporal scales, from a season to hundreds of thousands and even millions of years. In addition, EMICs include, although often in parameterised form, most of the processes described in GCMs (Claussen et al., 2002). On the other hand, they are efficient enough to resolve climate dynamics on the event-scale (kyr) and are thus especially useful for the study of paleoclimate dynamics since they allow exploring the complex behaviour of Earth's climate system as an integrated multi-component system with non-linearly coupled processes. Often EMICs incorporate more sub-component models (e.g. sediments) and climatic variables than GCMs (Petoukhov et al., 2005). Furthermore, they facilitate large ensemble experiments needed for climate sensitivity studies to external forcings (e.g. Dalan et al., 2005; Goodwin et al., 2009) and can provide guidance for more detailed investigations using more complex GCMs. Also, due to their fast computation, EMICs are able to integrate long-term processes like carbonate preservation in marine sediments (Ridgwell and Hargreaves, 2007; Tschumi et al., 2011) which is important for regulating atmospheric CO<sub>2</sub> concentrations on time-scales of thousands to tens of thousands of years (Ridgwell and Zeebe, 2005). The model design is generally driven by the underlying scientific questions, the considered temporal and spatial scales, as well as by the expertise of the research group. As a consequence, different components of the climate system are described with different levels of complexity. Most EMICs emerged from comprehensive, dynamic climate models that were used to study contemporary climate change and thus rely on sophisticated coupled ocean-atmosphere GCMs with integrated ice sheet dynamics (e.g. McGuffie and Henderson-Sellers, 2005). Terrestrial vegetation and weathering dynamics in EMICs are




Fig. 4. Timeline for EMIC and Earth system box model applications. Diamonds represent steady-state studies, solid lines display transient simulations for the specific time period and dotted lines steady-state studies for the respective period without a specified time. The size of the diamonds and thickness of the lines reflect the number of studies for this time period (also compare Table 5 and 6 for the list of model applications). The geological timescale abbreviations for the periods are: N: Neogene, Pg: Paleogene, K: Cretaceous, J: Jurassic, T: Triassic, P: Permian, C: Carboniferous, D: Devonian, S: Silurian, O: Ordovician, ε: Cambrian. Note the change of timescale!

often included on the basis of dynamic models (such as VECODE in CLIMBER-2; Brovkin et al., 2002a; and RoKGeM in GENIE; Colbourn et al., 2013, respectively). A major caveat of many existing models is the lack of a sophisticated sedimentary biogeochemical model for organic carbon (compare Section 3.3). Notable exceptions are Bern 3D which includes a vertically-integrated, dynamic model considering oxic degradation and denitrification of organic carbon (Tschumi et al., 2011); the box model MBM, integrating a vertically-resolved advection-diffusion-reaction model for solid and solute species (MEDUSA, Munhoven, 2007); and DCESS, using a semi-analytical, iterative approach considering (oxic and anoxic) organic matter remineralisation (Shaffer et al., 2008).

### 3. Overview of the primary processes involved in marine carbon cycling

Over the geological past, the intensity of the biological pump has probably experienced significant spatial and temporal variability that can be directly linked to perturbations of the global carbon cycle and climate (e.g. Sigman and Boyle, 2000; Kohfeld et al., 2005; John et al., 2014; Ma et al., 2014). Therefore, the description of the biological carbon pump exerts an important control on the performance and predictive abilities of Earth system models. Simulating the ocean's carbon pump is made difficult by the plethora of processes that govern the formation of particulate organic and inorganic carbon (POC, PIC) and dissolved organic carbon (DOC) in the euphotic zone, its export to depth and its subsequent degradation, dissolution or burial in the sediments. Unlike for ocean circulation, a fundamental set of first principle equations that govern nutrient cycling still remains to be defined, assuming they even exist. Thus a wide range of model approaches of varying sophistication have been put forward for specific problems (e.g. Bacastow and Maier-Reimer, 1990; Najjar et al., 1992; Maier-Reimer, 1993; Sarmiento et al., 1993; Six and Maier-Reimer, 1996). The following sections critically review important processes of the biological pump and their representation in paleoclimate models. They are separated into biological production in the surface ocean (Section 3.1), carbon dynamics in the deeper ocean (Section 3.2) and benthic processes (Section 3.3). The numerical representations are generally described moving from simple/empirical to more dynamic/mechanistic approaches. Even though the silica cycle is strongly interrelated with the biological pump (e.g. opal as a ballasting material; Klaas and Archer, 2002) we limit our review to carbon containing compounds and refer the interested reader to Ragueneau et al. (2000) and Tréguer and Rocha (2013) for reviews on the ocean silica cycle.

#### 3.1. Biological production

Biological production is controlled by the availability of light, nutrients, and trace metals, as well as phytoplankton speciation, temperature, and grazing. At steady state, nutrients removed from surface waters in the form of descending particulate matter are balanced by the upward advective and diffusive supply of dissolved nutrients that support new production in surface waters (Eppley and Peterson, 1979). The production of PIC, POC and DOC in the surface ocean is thus driven by complex recycling and transformation pathways within the euphotic ecosystem (e.g. Sarmiento and Gruber, 2006). High data requirements, excessive computational demands, as well as the limited transferability of existing comprehensive ecosystem model approaches to the geological timescale compromise the application of these models in a paleo-context. Therefore, paleoclimate models generally treat surface ocean biogeochemical dynamics in a simplified way (see Fig. 5 for an overview).

##### 3.1.1. Organic carbon production

Model approaches can be broadly divided into two different classes (Fig. 5). A first class of models simply relates the whole-community

export production directly to the availability of nutrients within the euphotic zone using either a nutrient restoration scheme towards a fixed concentration distribution ( $NU^*$ , e.g. Najjar et al., 1992), or a Michaelis-Menten term for uptake kinetics (e.g. Maier-Reimer, 1993). These export production models sometimes account for the more refractory DOC pools by assuming that a fixed fraction,  $\delta$ , of the produced carbon is converted to DOC in the production zone. The second class of models explicitly resolves, although in a very reduced and simplified way, part of the biological complexity that drives carbon production in the euphotic zone (Six and Maier-Reimer, 1996; Mouchet and François, 1996; Keller et al., 2012). In these models, the export production is driven by a pool of phytoplankton whose growth is controlled by the availability of nutrients, light, as well as temperature. Upon death, organisms feed the fast sinking particulate organic carbon pool and POC export production is thus determined by grazing and mortality rates. The biological model of Mouchet and François (1996) applies a simple maximum grazing rate with a Michaelis-Menten term that allows for a non-linear closure of the system. Six and Maier-Reimer (1996) on the other hand explicitly resolve the dynamics of nutrients, phytoplankton, zooplankton and detritus (NPZD-model) in the euphotic layer and account for the production of fecal pellets by both carnivores and herbivores. Furthermore, their model assumes that DOC is produced by exudation from phytoplankton and zooplankton excretion.

All global carbon cycle box models integrate a simple export production model including highly parameterised maximum production terms, since their low spatial resolution does not permit the resolution of latitudinal temperature and light variations. Some EMICs also apply simple export production models, however, the maximum export production depends here on the ambient temperature and/or light conditions. The UVic model, CLIMBER-2 and LOVECLIM have more complex biological schemes, following Keller et al. (2012), Six and Maier-Reimer (1996) and Mouchet and François (1996), respectively. Although there is still a controversy about nitrogen or phosphorus being the ultimate limiter of oceanic primary production at geological time-scales (Smith, 1984; Falkowski et al., 1998; Tyrrell, 1999), most paleoclimate models estimate export production directly from available surface phosphate concentrations, implicitly assuming that nitrogen fixation compensates for a potential nitrogen limitation.

##### 3.1.2. Pelagic calcium carbonate production

Today, the surface ocean is largely oversaturated with respect to carbonate phases (see Box 1 for a carbonate primer; Ridgwell and Zeebe, 2005). Nevertheless, due to the kinetically unfavourable initial step of crystal nucleation, abiotic carbonate precipitation is rare and only occurs in extreme environments as cements or ooids (e.g. Morse and He, 1993; Schneider et al., 2006). In the open ocean, the most important groups for the production of calcium carbonate ( $CaCO_3$ ) are marine organisms like coccolithophores, foraminifera and pteropods. These three groups are responsible for up to 70% of global  $CaCO_3$  precipitation (Milliman and Droxler, 1996). Coccolithophores are a group of algae (mostly unicellular) that form an outer sphere of small calcite crystals, known as coccoliths. The other two groups are heterotrophic zooplankton. Foraminifera form skeletons made out of calcite and pteropods, a general term for a group of molluscs, produce crystals made out of aragonite (the thermodynamically less stable form of calcium carbonate).

As these biological processes and their links to ocean geochemistry are very complex, most paleoclimate models calculate marine carbonate export production as a fixed fraction of POC export production (the  $r_{PIC/POC}$  "rain ratio", compare Fig. 5 and e.g. Yamanaka and Tajika, 1996). However, there is no simple means of deriving the value of  $r_{PIC/POC}$  as it is highly dependent on the local ecosystem structure (Shaffer, 1993). Paleoclimate models that integrate biological models generally apply a formulation that relates the rain ratio to the ecosystem structure. Otherwise, highly parameterised formulations on the basis of ambient temperature conditions (e.g. Maier-Reimer, 1993; Heinze





Fig. 5. Overview of model approaches that are applied in paleoclimate models to calculate surface ocean production/export production.

et al., 1999; Archer et al., 2000b; Sarmiento et al., 2002) or the saturation state with respect to calcite are used (Ridgwell et al., 2007; Arndt et al., 2011). Only a small number of paleoclimate models distinguish between calcite and aragonite fractions as most of the global PIC export is driven by low-Mg calcite from coccolithophores and foraminifera (Iglesias-Rodriguez et al., 2002; Morse et al., 2007). The aragonite cycle is still poorly quantified and therefore, its rain ratio is considered in a few models only: LOCH/LOVECLIM (Mouchet and François, 1996), MBM (Munhoven and François, 1996; Munhoven, 2007), and optionally in Bern 3D (Gangstø et al., 2011). Estimates for the fraction of aragonitic pteropods or heteropods in the global ocean

carbonate rain are in the range of 10–50% (Bernier, 1977; Bernier and Honjo, 1981; Fabry, 1989).

### 3.2. Carbon (re)cycling in the ocean interior

The remineralisation of biogenic particles exported from the surface ocean redistributes carbon and nutrients in the ocean exerting an important influence on atmospheric CO<sub>2</sub> levels. Changes in the depth-distribution of POC remineralisation controls the sequestration of CO<sub>2</sub> in the water column on time-scales of decades to centuries (Kwon et al., 2009) whereas changes in the ratio of POC to CaCO<sub>3</sub> in fluxes reaching



**Fig. 6.** Overview of model approaches that are applied in paleoclimate models to calculate the depth profiles of POC fluxes in the water column.

the sediment may change  $\text{CO}_2$  over thousands to millions of years (Archer and Maier-Reimer, 1994; Roth et al., 2014). However, despite considerable effort to identify potential processes that control the fate of organic carbon in water column (e.g., Joint Global Ocean Flux Study (JGOFS, Buesseler and Boyd, 2009); VERTICAL Transport In the Global Ocean (VERTIGO, Trull et al., 2008)), the key processes are still not well understood (Boyd and Trull, 2007).

### 3.2.1. Particulate organic carbon

**3.2.1.1. Observations.** Field observations across the global ocean show that a large fraction of particulate organic carbon is efficiently degraded by bacterial activity as it sinks through the water column and that the recycling process is exponentially faster in the mesopelagic layers of the ocean than in the deeper ocean (e.g. Martin et al., 1987; Lutz et al., 2002; Honjo et al., 2008, compare Figs. 2 and 7). This is generally explained with either increasing particle sinking speeds with depths (Berelson, 2001) or a decrease in organic matter reactivity with depth/

age because as POC sinks, labile organic components are degraded more readily, leaving more refractory material behind (e.g. Dauwe et al., 1999). However, the overall importance of this so-called selective preservation remains disputed (e.g. Hedges et al., 2001; Gupta et al., 2007), and little is known about the mechanistic control on decreasing organic matter degradability during sinking. Other mechanisms, such as physical protection (e.g. Keil and Hedges, 1993) likely also contribute to decreasing organic matter reactivity. Compilations of globally distributed sediment trap data show that the rate of attenuation with depth varies generally with latitude. However, very little is known about what controls this geographical variability and different observations are seemingly inconsistent (see e.g. Henson et al., 2012; Marsay et al., 2015). Various mechanisms have been postulated to explain these spatial patterns, focusing again on either changes in the sinking rate or the degradation rate of POC.

As the density of typical organic matter ( $\sim 1.05 \text{ g cm}^{-3}$ , Logan and Hunt (1987)) is close to that of seawater ( $\sim 1.02\text{--}1.03 \text{ g cm}^{-3}$ ) POC




Fig. 7. Comparison of discussed POC flux representations and global POC flux data from Lutz et al. (2002). Observations shallower than 1.5 km have been divided by 0.4 to account for the potential undertrapping error (open symbols). Observations: ■/□: Atlantic Ocean, ◆/◇: Pacific Ocean, +/◊: Indian Ocean, ▲/△: Greenland Sea. POC flux data has been normalised to regional POC export estimates given in Lutz et al. (2002) - Table 2.

needs to aggregate or a source of weight in order to contribute significantly to the organic matter flux to the deep ocean. Strong global correlations between inorganic minerals (such as  $\text{CaCO}_3$ , opal and lithogenic material) and POC fluxes in the deep ocean led to the hypothesis that the denser minerals increased the density, and therefore sinking rate, of POC (the “ballast hypothesis”) (Armstrong et al., 2001; Klaas and Archer, 2002). Deep ocean POC fluxes are thus driven by the local biomineral (i.e. high in calcite-dominated and low in opal-dominated regions.) However, analyses of these relationships on temporal (Lam et al., 2011) and spatial scales (Wilson et al., 2012) have questioned the relative role of minerals as ballast material. The correlations may also reflect other processes governing the association between POC and minerals such as the presence of transparent exopolymer particles (TEP) that stimulate aggregation Passow (2004), De La Rocha et al. (2008). Another explanation for the observed regional POC flux variability is a temperature dependence for the degradation of organic matter. Temperature is a primary determinant for bacterial degradation rates (Gillooly et al., 2001; Matsumoto, 2007) and it has been shown that degradation rates are more sensitive to temperature than photosynthesis (López-Urrutia et al., 2006; Regaudie-de Gioux and Duarte, 2012). Thus, warmer waters are characterised by faster degradation rates and therefore shallower remineralisation (Marsay et al., 2015).

A hypothesis that combines elements of the previous mechanisms specifies ecosystem structure and specifically the extent of recycling of organic matter in the euphotic zone as an important factor governing the observed spatial variability in POC flux (Henson et al., 2012; Le Moigne et al., 2012, 2014). The reasoning is that in high export production areas (generally colder high latitudes) aggregates are rather fresh and loosely packed as they are a result of strong, seasonal diatom blooms, making them prone to rapid degradation. In low export production areas (warmer low latitudes) the material being finally exported has been processed multiple times in the euphotic layer and is therefore tightly-packed, highly refractory and thus experiences reduced microbial degradation at mesopelagic depths (Henson et al., 2012).

An increasing number of mechanisms have been identified that may contribute to changes in the attenuation of POC flux in the water column. However, as highlighted above, these mechanisms are potentially interlinked and difficult to distinguish from current observations. This represents a significant source of uncertainty surrounding the magnitude and sign of ocean carbon cycle feedbacks to changes in atmospheric  $\text{CO}_2$  and climate (Barker et al., 2003; Riebesell et al., 2009).

**3.2.1.2. Numerical approaches.** Most Earth system models impose static POC depth profiles and are therefore based on simple fitting exercises to limited data sets rather than on a mechanistic understanding of the underlying processes. This is surprising, since more sophisticated models have long been used to study organic carbon degradation dynamics in soils and marine sediments over different time-scales (e.g. Arndt et al., 2013). Even though still far from providing an appropriate representation of the plethora of different mechanisms that control organic carbon degradation, these models have proven useful in describing the degradation dynamics of organic carbon from different sources, in different environments, under changing redox-conditions and over time-scales (e.g. Arndt et al., 2013). The applied approaches in ocean biogeochemical models can be broadly divided into two groups: the primarily empirical approaches and more mechanistic approaches of various levels of complexity.

In the early 1980’s net primary productivity (NPP) in the surface waters was assumed to be most important for vertical POC flux. The empirical algorithm of Suess (1980) (Eq.(1) in Fig. 6) describes POC flux to depth as a function of NPP, scaled to depth below the sea-surface. However, Bishop (1989) demonstrated that export (or new) production of POC (i.e. POC exported from the euphotic zone) predicts POC flux in deeper waters more reliably and is used until now in biogeochemical models. Other empirically derived relationships used in paleoclimate models are intentionally or unintentionally related to more mechanistic descriptions of organic carbon degradation. Power-law (Martin et al., 1987), single exponential (Volk and Hoffert, 1985; Shaffer, 1996; Heinze et al., 1999) or double exponential (Lutz et al., 2002; Andersson et al., 2004) relationships are applied in most of the paleoclimate models (Table 1). Although these models are usually empirical as well, they can be directly derived from the kinetic first-order rate law of organic matter degradation:

$$\frac{d\text{POC}}{dt} = k \cdot \text{POC} \quad (1)$$

and the chosen function reflects certain assumptions about the organic matter pool and its degradation rate  $k$ . Therefore, the representation may be more directly related to some underlying bioenergetic drivers (Berner, 1980b; Boudreau, 1997). The rate constant of organic carbon degradation,  $k$ , is usually interpreted as a measure of the reactivity of the macromolecular organic matter towards hydrolytic enzymes and is thus assumed to primarily depend on the macromolecular composition of organic matter. It therefore encompasses not only the original composition of the exported organic carbon, but also its evolution during sinking (Arndt et al., 2013). The simplest form of this approach assumes that the organic carbon constitutes one single pool, which is degraded at a constant rate. This approach is equivalent to the so-called 1G-Model of organic matter degradation (Berner, 1964) that has been widely used in diagenetic modelling (Boudreau, 1997). Its steady-state solution is given by a simple exponential decrease of organic carbon flux with depth that is controlled by the reactivity of organic carbon,  $k$ , and the settling velocity,  $w$  (Eq.(2) in Fig. 6; see Sarmiento and Gruber, 2006 for a derivation). Yet, this simple exponential model represents merely a linear approximation of the complex degradation dynamics and the first-order degradation constant,  $k$ , represents a mean value for the heterogeneous mixture of organic compounds. It should be noted that such simplification is reasonable only if the degradability of different compounds does not vary by more than one order of magnitude (Arndt et al., 2013).

Under the assumption that the organic carbon degradation rate decreases linearly with depth, a power-law functionality for POC flux can be derived from kinetic first-order principles (compare supporting information of Lam et al., 2011). Most commonly used is the description of Martin et al. (1987) – Eq.(3) in Fig. 6. Martin et al. (1987) fitted a number of sediment trap POC flux measurements from six different locations in the northeast Pacific Ocean to a simple power-law. The expression scales deep fluxes to POC export from the euphotic zone

**Box 2**

The paleoclimate model GENIE and the experiment set-up.

**The “Grid ENabled Integrated Earth system model” (GENIE)**

The basis GENIE is a 3D-ocean circulation model coupled to a fast energy-moisture balance 2D-atmosphere model (“C-GOLDSTEIN”, Edwards and Marsh, 2005). To help understand the oceanic carbon cycle and its role in regulating atmospheric CO<sub>2</sub> concentrations the model has been extended with a ocean biogeochemistry representation for a variety of elements and isotopes (Ridgwell et al., 2007). The ocean model is implemented on a 36 × 36 equal-area horizontal grid and 16 z-coordinate levels in the vertical. Despite its lower resolution, GENIE is able to reproduce the main nutrient and dissolved inorganic carbon δ<sup>13</sup>C features of the modern ocean (Holden et al., 2013; Ridgwell et al., 2007). The same model physics have been applied before to an early Eocene and Late Cretaceous bathymetry and continental configuration and successfully modelled various oceanic and sedimentary properties related to the biological carbon pump (John et al., 2014; Kirtland Turner and Ridgwell, 2013; Monteiro et al., 2012; Ridgwell and Schmidt, 2010).

**Experiment set-up: oceanic POC-flux representations (Figs. 8, 9 and 10)**

The GENIE set-up for the modern and Eocene experiments is identical to John et al. (2014). The OAE2 experiments adopt the carbon cycle boundary conditions of Monteiro et al. (2012) using 2 times modern oceanic phosphate concentration (4.5 μmol PO<sub>4</sub> kg<sup>-1</sup>) and 4 times preindustrial atmospheric CO<sub>2</sub> (1112 ppmv). However, we do not consider nitrogen in our simulations, thus phosphate being the only productivity limiting nutrient. All experiments are run for 10,000 years in order to equilibrate the ocean biogeochemistry. In the reference model set-up, POC-flux throughout the water-column is modelled using the fixed double-exponential approach (Eq.(4), Fig. 6). The temperature-dependent remineralisation uses the formulation as discussed for Fig. 7. The free parameters of this approach (i.e. two rate constants (A) and the initial refractory fraction of POC) were calibrated for the modern ocean to find the best fit with the double-exponential approach for the global POC-flux profile. The two approaches are then used under a preindustrial (modern) configuration (Cao et al., 2009), an early Eocene set-up (Ridgwell and Schmidt, 2010) and an OAE2 configuration (Monteiro et al., 2012).

**Experiment set-up: sediment representations (Figs. 16 and 17)**

For this series of experiments GENIE employs the fixed double-exponential POC-flux scheme but is configured with two different sediment boundaries for the modern, Eocene and OAE2 worlds: first, the so-called reflective boundary, where essentially no sediment interactions take place and all particulate species reaching the seafloor are instantaneously remineralised to dissolved carbon and nutrients. The second boundary represents the other end of the spectrum and assumes that the entire deposition flux of POC is buried in the sediments (conservative or semi-reflective boundary). In order to calculate a steady-state situation we configure the model as a “closed” system where the burial loss to the sediments is balanced by an additional weathering input of solutes to the ocean through rivers. Because of the sediment interactions the experiments are run for 20,000 years to reach equilibrium.

( $z_{\text{eup}}$ ) and flux attenuation with depth is parameterised with a constant parameter  $b = 0.858 \pm 0.1$ . The majority of existing paleoclimate models integrate the Martin curve in its original form with a constant parameter  $b$  (Table 1) that does not change temporally or spatially. Its popularity mainly stems from the fact that a power-law represents a mathematically simple way to describe the sharp decrease of organic carbon fluxes in mesopelagic layers, while still maintaining a flux at depth.

Yet, in reality, the organic carbon flux to depth is composed of many specific and very heterogeneous compounds with a continuum of individual decay rates, distributed from very labile compounds that are degraded within several hours or days to highly condensed compounds that persist for hundreds of thousands or even millions of years. The assumption of a single organic carbon pool that is degraded with a constant or linear degradation rate is thus not consistent with natural conditions. The bulk POC flux can be subdivided into a number of compound classes that are characterised by different degradabilities  $k_i$ . This approach follows the multi-G approach proposed by Jørgensen (1978) and Berner (1980a) to describe organic carbon degradation in marine sediments. The rapid depletion of the more reactive compound class results in a decrease of total organic carbon degradability with water depth and therefore provides a more realistic representation of organic carbon degradation dynamics (Bishop, 1989; Wakeham et al., 1997). Lutz et al. (2002) propose to describe organic carbon flux to depth using a double exponential equation including degradation length scales for a labile and a refractory fraction of POC (Eq.(4) in Fig. 6). For instance GENIE integrates such a double exponential expression that can capture the rapid flux attenuation in subsurface waters as well as the slower flux attenuation in deep waters. In theory, this 2G-Model could be expanded by introducing other POC compound classes, but as they are difficult to identify from observational data their number is generally restricted to three (Jørgensen, 1978; Middelburg, 1989).

An alternative to these models are Reactive Continuum Models (RCMs) of organic matter degradation (Eq.(5) in Fig. 6). These models assume a continuous distribution of reactive types, thus avoiding the highly subjective partitioning of POC into a limited number of compound groups (e.g. Middelburg, 1989; Boudreau and Ruddick, 1991). A newer version of GEOCLIM (GEOCLIM *reloaded*; Arndt et al., 2011) applies the RCM to quantify POC fluxes below the euphotic zone. The RCM provides a direct conceptual link between the composition of organic matter and its degradability. The total amount of organic matter is represented as the integral of the distribution function of reactive types over the entire range of possible degradation rate constants. Each member is degraded according to a first order rate law. The RCM thus explicitly integrates the effect of compound-specific reactivities on organic matter degradation and captures the decrease in organic matter degradability with depth/age. However, the RCM approach requires the determination of realistic differential reactivities of specific components rather than an average reactivity for the total POC pool which can be challenging.

**3.2.1.3. Comparing numerical approaches.** Fig. 7 compares some of the discussed approaches with a compilation of globally distributed POC flux observations from the modern day (Lutz et al., 2002). The Henson approach uses the Martin equation but calculates the  $b$ -value in relation to local SST ( $b = (0.024 \cdot \text{SST}) - 1.06$ , compare Henson et al. (2012, Fig. 4f)). A mean SST of 17.4 °C leads to a global mean  $b$ -value of  $-0.643$  and therefore predicts higher residual POC flux in the deep ocean as the Martin model, showing how sensitive this approach is to changing  $b$ -values. The ballasting-method assumes that part of the POC export is associated with a ballasting mineral (here calcium carbonate) and the excess POC is considered the labile fraction (Armstrong et al., 2001; Klaas and Archer, 2002):

$$F_{\text{POC}}(z) = \lambda_{\text{Ca}} \cdot F_{\text{Ca}}(z) + F_{\text{labile}}(z), \quad (2)$$




Fig. 8. Modern and Eocene global profiles for a GENIE set-up using a double-exponential and temperature-dependent POC remineralisation approach. See Box 2 for more information about the experiment set-up.

where  $F_{\text{POC}}(z)$  is the total POC flux at depth  $z$ ,  $\lambda_{\text{Ca}} = 0.126$  the carrying coefficient of  $\text{CaCO}_3$ ,  $F_{\text{Ca}}(z)$  and  $F_{\text{labile}}(z)$  the total calcium carbonate and labile POC mass flux as calculated with GENIE. The temperature dependent remineralisation follows the description of John et al. (2014) using an Arrhenius-type equation to predict the temperature dependent remineralisation rate  $k(T)$ :

$$k(T) = A \cdot \exp\left(-\frac{E_a}{RT}\right), \quad (3)$$

where  $R$  is the gas constant and  $T$  the absolute temperature. To distinguish between labile and refractory POC two activation energies  $E_a$  (55 and 80  $\text{kJ mol}^{-1}$ ) and two rate constants  $A$  ( $9 \times 10^{11}$  and  $1 \times 10^{14} \text{ year}^{-1}$ ) were chosen (as calibrated in John et al., 2014). The

mean temperature profile for the global ocean has been taken from the World Ocean Atlas (Locarnini et al., 2013). In principle, all shown formulations are able to capture the characteristic POC-flux profiles for the modern day ocean (Fig. 7). However, as the parameters applied in paleoclimate models are constrained to modern-day observations (e.g. Martin et al., 1987; Honjo et al., 2008) and the performance of the models strongly depend on the parameter choice, their predictive ability under different climatic conditions is seriously compromised.

To illustrate the impact of using a static versus a mechanistic POC-flux representations on ocean biogeochemistry we compare the fixed double-exponential with the temperature-dependent remineralisation approach using the paleoclimate model GENIE (see Box 2 for more information on the model and experiment set-up). Fig. 8 compares




Fig. 9. Anomaly plots (temperature-dependent minus double-exponential POC remineralisation approach) for the modern (left panel) and the Eocene (right panel). Top: export production (i.e. POC flux at 40 m). Bottom: depth integrated remineralisation concentration of  $\text{PO}_4$ .

Table 3  
Characterisation of major DOC fractions (after Hansell, 2013).

Fraction	Inventory	Removal rate	Lifetime	Main occurrence
Labile	(Pg C) < 0.2	( $\mu\text{mol C kg}^{-1} \text{ year}^{-1}$ ) ~100	Hours to days	Directly recycled
Semi-labile	$6 \pm 2$	~2–9	Months to years	Upper mesopelagic zone (< 500 m)
Semi-refractory	$14 \pm 2$	~0.2–0.9	Years to decades	Mesopelagic zone (< 1000 m)
Refractory	$642 \pm 32$	~0.003	Centuries to millennia	Everywhere

**Table 4**

DOC representation in paleoclimate EMICs. Note, none of the reviewed global carbon cycle box models represents a DOC pool.


Model	DOC fraction	Lifetime	Reference
Bern 2.5D	Constant inventory	Dynamic	Marchal et al. (1998b)
Bern 3D	Constant inventory	Dynamic	Marchal et al. (1998b)
CLIMBER-2	Labile/semi-labile	Days to 1 year	Six and Maier-Reimer (1996)
GENIE	Semi-refractory	40 years	Brovkin et al. (2002a)
	Semi-labile	0.5 years	Ridgwell et al. (2007)
	Semi-refractory	20 years	Ma and Tian (2014)
	Refractory	10,000 years	Ma and Tian (2014)
GEOCLIM	–	–	–
LOVECLIM	Semi-refractory	20–22 years	Goosse et al. (2010)
MESMO	Semi-labile	0.5 years	Ridgwell et al. (2007)
MoBidiC	Semi-refractory	8.6 years	Crucifix (2005)
UVic	Semi-labile/semi-refractory	2–6 years	Schmittner et al. (2005)

global POC, oxygen and nutrient profiles for a modern and a warm Eocene (55.5–33.7 Ma) climate. The global POC flux profile (normalised to export flux) for the static double-exponential just changes slightly in the Eocene experiment, whereas the temperature-dependent profile gets significantly shallower and a smaller POC fraction reaches the sediments due to warmer ocean temperatures (Fig. 8A). The global O<sub>2</sub> profile for the temperature-dependent parameterisation in the Eocene also shows this shoaling compared with the modern (Fig. 8B). But the temperature dependence of POC remineralisation has also major impacts on the modern ocean as can be inferred from the global O<sub>2</sub> profile, showing a different shape compared with the double-exponential scenario. However, in contrast the global profiles for PO<sub>4</sub> do not change significantly (Fig. 8C), a result that can be attributed to the fixed initial global phosphate inventory imposed onto the model (compare also, e.g., Kriest and Oschlies, 2013). The POC remineralisation scheme has not just significant impacts on global mean biogeochemical values but also affects their spatial distribution. Fig. 9 shows anomaly plots (temperature-dependent minus double-exponential) for export production (A and B) and depth integrated remineralisation concentrations of PO<sub>4</sub> (C and D). The differences for modern conditions are marginal (Fig. 9, A and C) as the temperature-dependent POC flux has been calibrated to the double-exponential flux

for these conditions. However, for the Eocene, the temperature-dependent export production increases globally (9.6 Pg C year<sup>-1</sup> compared to 5.9 Pg C year<sup>-1</sup> in the double-exponential simulation), in particular in warm equatorial regions (Fig. 9B). Also the depth integrated PO<sub>4</sub> remineralisation is significantly higher (Fig. 9D). Both can be explained by the shallower, temperature dependent degradation of POC in the warmer Eocene, leading to higher PO<sub>4</sub> availability in the upper ocean and therefore an absolute increase in global productivity.

Fig. 10 compares the two remineralisation approaches for another extreme event in Earth history, the Late Cretaceous oceanic anoxic event (OAE2). Shown are modelled H<sub>2</sub>S concentrations, another indicator for ocean redox conditions, for the two GENIE configurations for OAE2. Euxinia is defined by the occurrence of free hydrogen sulphide in the water column, which is characteristic of anoxia as H<sub>2</sub>S is produced by sulfate reduction when oxygen is depleted. Fig. 10A + B highlights the significant impact the POC-flux representation has on photic zone euxinia, with far more H<sub>2</sub>S predicted when the temperature-dependent approach is used, a result that can be explained with warmer ocean temperatures leading to more POC degradation in the upper ocean (POC profile for OAE2 is similar to the Eocene Fig. 8, not shown here). However, as the majority of POC is degraded in the upper 500 m when using the temperature-dependent approach, less H<sub>2</sub>S is produced in the deeper ocean (Fig. 10C + D).

**3.2.1.4. Summary.** The different model results highlight that the current lack of a mechanistic theoretical framework to model POC flux seriously compromises the applicability of paleoclimate models to extreme and rapidly changing environmental conditions. A complex process interplay controls organic carbon degradation in the Earth system. It has been shown that the availability of oxygen, variable redox-conditions, euxinic environmental conditions or changing organic carbon sources can exert profound impact on organic matter reactivity and thus degradation and burial (e.g. Demaison and Moore, 1980; Canfield, 1994; Sinninghe Damsté et al., 1998; Meyers, 2007; Arndt et al., 2009). A robust mechanistic framework is therefore a crucial prerequisite to increase the predictive ability of POC flux models, especially under changing environmental conditions that typically characterise past carbon-cycle perturbations.



**Fig. 10.** Model comparison of H<sub>2</sub>S concentration (i.e. euxinia) during OAE2 for a GENIE configuration with double-exponential (left) and temperature-dependent (right) POC remineralisation approach. (A + B): photic zone euxinia showing modelled H<sub>2</sub>S concentration at 80–200 m. (C + D): vertical profile of H<sub>2</sub>S in the East Pacific Ocean (–90° longitude, area indicated by black rectangles in A + B).




Fig. 11. Overview of model approaches that are applied in paleoclimate models to calculate the depth profiles of PIC fluxes.

### 3.2.2. Dissolved organic carbon

**3.2.2.1. Observations.** With a size of about 662 Pg C, marine dissolved organic carbon (DOC) is comparable to the amount of carbon in the atmosphere (Hansell and Carlson, 2014). Recently, various studies have re-emphasized the importance of DOC in the ocean and its contribution to the biological pump (e.g. Hansell et al., 2009; Jiao et al., 2010; Hansell, 2013). Most marine DOC is produced (together with POC and PIC) by phytoplankton in the surface ocean and accounts for about 30–50% of the primary production (Ducklow et al., 1995; Biddanda and Benner, 1997). A large part of the produced DOC belongs to the labile or semi-labile DOC pool. The labile fraction is directly recycled in the euphotic zone (lifetime hours to days), whereas the semi-labile DOC mostly consists of carbohydrates that are degraded by heterotrophic processes in the upper 500 m of the ocean (see Table 3). Therefore, semi-labile DOC represents an important contributor to the biological carbon pump. Due to short lifetimes the labile and semi-labile DOC fractions account for a mere 1% of the total DOC inventory and have a limited importance for longer carbon sequestration (Hansell, 2013). The remainder is transported further to deeper waters through the overturning circulation of the ocean or scavenging on sinking aggregates and can be broadly divided into a semi-refractory and refractory pool (Hansell et al., 2012). Semi-refractory DOC is mainly found at mesopelagic depth of the ocean (< 1000 m) and accounts for about 2% of the DOC inventory (lifetime years to multiple decades, Hansell et al., 2012). Hansell (2013) argues that most of the deep ocean DOC is largely unreactive and degrades on time-scales of several

hundreds to thousands of years. This refractory DOC pool thus survives several cycles of ocean overturning and represents the largest fraction of the total marine DOC reservoir (about 97%, Table 3). Therefore, it is mostly argued that semi-labile DOC largely dominates the upper ocean (< 500 m), while semi-refractory and refractory DOC represents most of the DOC in the deep ocean (Hansell, 2013). In contrast, Jannasch (1994) and Arrieta et al. (2015) hypothesise that most of the deep ocean DOC is in fact labile but that its very low concentrations limit their microbial utilisation. This “dilution hypothesis” is supported by the fact that most of the refractory DOC is still unclassified (Kujawinski, 2011) and little evidence exists to prove that it should not be available for microbial degradation.

**3.2.2.2. Numerical approaches.** Despite these uncertainties and its unquantified importance for the biological pump, none of the paleoclimate box models integrate an explicit description of DOC (Table 4). Most other models describe the heterotrophic degradation of just one single DOC pool by a first order degradation rate law:

$$\frac{\partial \text{DOC}}{\partial t} = k_{\text{DOC}} \cdot \text{DOC} \quad (4)$$

The Bern models, for instance, assume a constant oceanic DOC inventory (Marchal et al., 1998b). As a consequence, DOC is degraded with a first order degradation rate constant,  $k_{\text{DOC}}$ , that is dynamically adjusted in a way that DOC production in the euphotic zone is balanced by its degradation in the deep ocean. This approach has been first

proposed by Najjar et al. (1992) who argue that the rate constant for DOC degradation cannot be constrained on the basis of data available at that time. The DOC lifetime in MoBidiC has been calibrated to 8.6 years, whereas LOVECLIM applies a degradation rate constant depending on oxygen availability resulting in a DOC lifetime between 20 and 22 years. Both models therefore capture the dynamics of the semi-refractory DOC pool. The standard setup of GENIE accounts for a semi-labile DOC fraction with a lifetime of 0.5 years. GENIE also has the option to represent a second DOC pool which has been used by Ma and Tian (2014) to model a semi-refractory and a refractory DOC pool (lifetimes of 20 and 10,000 years) for the Last Glacial Maximum. Dissolved organic carbon in the CLIMBER family is simulated with the biological model of Six and Maier-Reimer (1996). Therefore, the concentration of DOC depends on exudation from phytoplankton, excretion from zooplankton and a nutrient dependent degradation constant (resulting in a lifetime between days and one year). The CLIMBER-2 model also provides the possibility to allocate 10% of the produced detritus to another, semi-refractory DOC pool with a lifetime of 40 years (Brovkin et al., 2002a). Paleoclimate models thus mainly account for the cycling of, what is operationally classified as, the semi-labile and semi-refractory DOC fractions and generally ignore the largest DOC reservoir and its contribution to the biological carbon pump via carbon sequestration.

**3.2.2.3. Summary.** A more realistic representation of DOC in paleoclimate models should include several fractions of DOC with different lifetimes (Kirchman et al., 1993) and the advantages of using a reactive continuum model for DOC should be tested. The lack of a better representation of DOC is often attributed to the limited knowledge about the processes and mechanisms involved in the generation and degradation of this carbon reservoir (e.g. Yamanaka and Tajika, 1997). Yet, over the recent years, considerable progress has been made in advancing our understanding of the ocean's DOC reservoir (e.g. Hansell and Carlson, 2014). In addition, a number of authors have recently argued that especially the refractory DOC pool may have played a central role for past carbon isotope excursions and associated climate change (e.g. Tziperman et al., 2011; Sexton et al., 2011). Sexton et al. (2011) suggested that an anoxic and stratified Eocene deep ocean may have facilitated the accumulation of a large refractory DOC reservoir. Periodic release and oxidation of this surficial carbon pool (about 1600 Pg C) as a consequence of changes in ocean circulation could explain the observed rapid decline in the  $\delta^{13}\text{C}$  record and associated climate warming. The rapid recovery of the global carbon cycle is for Sexton et al. (2011) an indicator that  $\text{CO}_2$  was sequestered again by the ocean and not by the slower process of silicate rock weathering. However, one problem with this hypothesis is the unknown sensitivity of DOC degradation to ocean oxygenation (compare e.g. Ridgwell and Arndt, 2014). Elucidating the role of DOC in general and the refractory DOC pool in particular for past carbon cycle and climate perturbations will thus require a better integration of DOC in paleoclimate models.

### 3.2.3. Particulate inorganic carbon

**3.2.3.1. Observations.** The cycling of particulate inorganic carbon (i.e.  $\text{CaCO}_3$ ) in the ocean also affects the biological pump and therefore atmospheric  $\text{CO}_2$ , but by more indirect mechanisms. Whether carbonates precipitate or dissolve can be directly linked to the saturation state ( $\Omega$ ) of the ocean (readers are referred to Box 1 for a brief primer on carbonate thermodynamics). Compared to the 5% of POC that is exported from the euphotic zone and reaching the sediments, a significantly higher amount of PIC is vertically transported to the bottom of the ocean (about 50% of the PIC export flux, Sarmiento and Gruber, 2006). The role of sinking particulate inorganic carbon in the biological pump is complex, because the deep dissolution of PIC is largely controlled by the degradation of sinking POC and releases alkalinity, which in turn titrates part of the  $\text{CO}_2$

released during POC degradation. In addition its high specific gravity plays a key role for the sinking rates of biogenic aggregates (the ballasting effect) and thus the residence time of particulate carbon in the ocean (e.g., Francois et al., 2002; Klaas and Archer, 2002). The mechanisms responsible for carbonate dissolution in the ocean are still matter of debate (Morse et al., 2007). Global observations showing that the depth of the lysocline coincides with the saturation horizon (e.g. Sarmiento and Gruber, 2006) have been used to imply that thermodynamic constraints are a dominant control on calcium carbonate preservation. However, a kinetic control on carbonate dissolution has been highlighted by in-situ experiments in the North Pacific (e.g., Berger, 1967) and laboratory studies reveal that dissolution rates increase in undersaturated waters (Morse and Berner, 1972; Berner and Morse, 1974). In addition, observational evidence even points to a partial dissolution of sinking carbonate above the saturation horizon (e.g. Milliman, 1993). Millero (2007) estimates, using global production estimates of  $\text{CaCO}_3$  and globally averaged deep water sediment trap data, that probably 40–80% of the calcium carbonate produced in the surface ocean dissolves in the upper water column. However, the mechanisms that drive the dissolution of carbonates above the lysocline remain enigmatic. The dissolution of carbonates within acidic micro-environments, such as the digestive system of zooplankton and marine aggregates has been evoked as an explanation for the shallow dissolution (e.g., Fiadeiro, 1980; Alldredge and Cohen, 1987; Milliman et al., 1999; Jansen et al., 2002), but no clear conclusions could be established. Alternative explanations involve more soluble forms of carbonates, such as aragonite and high-magnesium calcites (e.g., Iglesias-Rodriguez et al., 2002; Feely et al., 2004). In summary, the dissolution of carbonates in the ocean is much more dynamic than our present understanding is able to explain.

**3.2.3.2. Numerical approaches.** This complexity is generally not reflected in the existing paleoclimate models. In fact, the applied approaches reflect much of the uncertainty that exists about the main drivers of calcium carbonate dissolution in the global ocean. In general, two very different approaches for the simulation of PIC depth profiles are applied in paleoclimate models (Fig. 11). The majority of paleo-models simply assume an exponential decrease of the PIC flux below the euphotic layer. The applied degradation length scales are typically chosen to be broadly consistent with the PIC flux ratio inferred from observations by Anderson and Sarmiento (1994), as well as Tsunogai and Noriki (1991) and fall within the range between 1000 m and 3000 m. The GENIE model assumes that a fixed fraction of the PIC export production reaches the deep ocean, while only the remaining fraction is subject to an exponential decrease with water depth. The magnitude of the respective fractions as well as the length scale are chosen consistent with the general conclusions of Milliman et al. (1999), and more specifically, with the sediment trap observations of Martin et al. (1993). In HAMOCC for instance (Maier-Reimer, 1993) the downward flux of  $\text{CaCO}_3$  is attenuated with a length scale of 2000 m. However, for some applications (e.g. Archer et al., 2000a), 30% of the carbonate export production was assumed to reach the sea-floor unchanged. In general, although able to represent calcium carbonate dissolution above the lysocline, the exponential model strongly simplifies calcium carbonate dynamics in the ocean. It is completely decoupled from the saturation state and organic carbon degradation dynamics, the most important drivers of calcium carbonate dissolution. Therefore, the exponential model cannot account for a dynamic response of the PIC flux profile to changing environmental conditions and its applicability to past extreme events is thus questionable. An alternative approach to the exponential model is based on the calcium carbonate reaction kinetics and is commonly used in diagenetic modelling. The overall process of calcium carbonate dissolution is a complex multi-step process. Although, the general rate law,  $R_{\text{diss}}$ , can be derived from thermodynamic and kinetic considerations (e.g. Lasaga, 1981), the most frequently employed rate law is empirically derived




Fig. 12. Schematic representation of the main early diagenetic processes and redox zonation in marine sediments. Typically sediments consist of several biogeochemical zones (left, as proposed by Berner, 1980b). Oxygen, and other species, diffuse from the water column into the upper sediment layer (blue arrows, the oxic zone). Deeper layers are suboxic or anoxic (sulfidic, methanic), and are characterised by different reactions in which for instance nitrate, manganese(IV), iron(III) or sulfate ions are reduced (and re-oxidised). Exact depths, however, vary strongly and increase from the shelf to the deep sea. The depth sequence of the dominant remineralisation reaction of organic matter (in white, AOM: Anaerobic Oxidation of Methane) is reflected in the vertical pore water profiles of its reactants and products (white, concentration scales are arbitrary). Carbonate reaching the deep-sea sediments may dissolve during early diagenesis if the bottom water is undersaturated or if porewater metabolic processes (primarily aerobic degradation) cause further undersaturation in the sedimentary porewater (right).

(Morse and Berner, 1972; Morse and Arvidson, 2002):

$$R_{\text{diss}} = \left( \frac{A}{V} k \right) \cdot (1 - \Omega)^n = k_{\text{PIC}} \cdot (1 - \Omega)^n \quad \text{if } \Omega < 1 \quad (5)$$

where  $A$  is the total surface area of the solid,  $V$  is the volume of solution,  $k_{\text{PIC}}$  is the rate constant and  $n$  is the apparent order of reaction. A synthesis of dissolution kinetics has shown that the apparent order of reaction varies between 1 and 4.5 (Keir, 1980; Hales and Emerson, 1997; Morse and Arvidson, 2002). Some paleoclimate models (such as LOVECLIM and GEOCLIM *reloaded*) use a simpler version of the thermodynamic model. Instead of resolving the depth distribution of dissolution rate, they assume that calcium carbonate does not dissolve in the water column but solely at the bottom of the ocean (i.e. mimicking surface sediment dissolution.). The thermodynamic approach accounts for the direct link between the organic and inorganic carbon cycle. In addition, it allows a dynamic response of the simulated PIC flux to changes in ocean chemistry and saturation state. Furthermore, its theoretical framework is based on a mechanistic understanding of the underlying thermodynamic and kinetic drivers which gives it the potential for feedback with changes in ocean chemistry. However, it does not account for the abundantly observed shallow carbonate dissolution in the modern ocean and might thus underestimate the extent of calcium carbonate dissolution in the water column. What is 'missing' in our understanding and hence models of the full details of the transfer of carbonate from the ocean's surface to underlying sediments is still not clear and hence the importance of this omission is difficult to quantify.

### 3.3. Carbon (re-)cycling at the ocean floor and controls on burial

At the seafloor, biogeochemical diagenetic processes are influenced by biogeochemical cycling in the overlying bottom water and the upper ocean. Dissolved species diffuse from the bottom water into the sediments and particulate material (such as organic matter, calcium carbonate and opal) rain down on the sediments and fuel biogeochemical reactions. Diagenesis transforms a substantial part of the deposited material (e.g. via remineralisation and/or dissolution) and the resulting products (e.g., DIC, nutrients) may return to the water column. As such, diagenetic processes are key components in the global carbon cycle as they trigger a delayed response to changes in ocean and atmosphere geochemistry and control the removal of carbon from the ocean reservoir (e.g. Arthur et al., 1988; Berner et al., 1989; Berner, 1990; Archer and Maier-Reimer, 1994; Mackenzie et al., 2004; Ridgwell and Zeebe, 2005).

In marine sediments, carbon is buried as organic carbon or carbonate minerals (Fig. 12). Ultimately, only a small fraction of the organic carbon (generally 10–20% of the deposited organic carbon or  $< \sim 0.5\%$  of the gross primary production) escapes remineralisation and is eventually buried in the sediment (e.g. Emerson and Hedges, 1988; Middelburg and Meysman, 2007; Burdige, 2007). However, organic carbon burial rates have been shown to vary significantly across different environments (e.g., Canfield, 1994) and through geological history (e.g., Arthur et al., 1985). The relative fraction of the deposited organic carbon that is ultimately buried in marine sediments can range from 0 to 100% (e.g., Canfield, 1994). A plethora of different mechanisms has been proposed to explain the observed patterns of organic carbon burial in marine sediments but their relative importance remains elusive (Demaison and Moore, 1980; Ibach, 1982; Pedersen and




Fig. 13. Calcite dissolution fluxes as a function of the degree of bottom water saturation,  $\Omega$ . The blue dashed line represents the dissolution that would occur due to bottom-water undersaturation only, in the absence of any metabolic  $\text{CO}_2$  (in analogy to the simplified thermodynamic model, Fig. 11). The red dashed line is the total carbonate dissolution (due to bottom-water undersaturation + organic carbon degradation) considering two organic carbon components and an average ocean-basin respiration rate. The two organic carbon component model represents a slower and faster decaying organic carbon pool in the sediments. Note that the scenario including the release of metabolic  $\text{CO}_2$  drives more carbonate dissolution for all saturation states.

Source: Modified from Hales (2003).

Calvert, 1990; Sageman et al., 2003).

It has been shown that on a global scale, only 10–15% of carbonate produced escapes dissolution and is buried in accumulating sediments (Milliman and Droxler, 1996; Archer, 1996b; Feely et al., 2004). Carbonate burial strongly depends on the saturation state of bottom- and porewaters. Because oceanic waters become increasingly less saturated at greater depth, deep-sea sediments are typically completely devoid of carbonate minerals. Furthermore, carbonate preservation is strongly influenced by the breakdown of organic matter and the release of metabolic  $\text{CO}_2$  (Archer, 1991; Hales, 2003; Ridgwell and Zeebe, 2005; Jourabchi et al., 2005). Fig. 13 illustrates potential differences in carbonate dissolution fluxes as a function of bottom-water saturation ( $\Omega$ ) for two different scenarios. The first considers only bottom-water undersaturation, whereas the second takes bottom-water undersaturation and the release of metabolic  $\text{CO}_2$  into account. The burial fluxes and efficiencies of carbonates and organic carbon are thus strongly influenced by early diagenetic processes, as well as their feedbacks on ocean biogeochemistry (Archer and Maier-Reimer, 1994; Mackenzie et al., 2004; Ridgwell and Hargreaves, 2007).

In marine sediments, geochemical processes are tightly coupled and geochemical species may undergo several recycling and transformation loops (e.g., authigenic mineral precipitation/dissolution) before they are either buried or diffuse back to the water column. This complex process interplay complicates the interpretation of the sedimentary record, one of the major climate archives on Earth. Coupled paleoclimate models, which include a mechanistic description of all the feedback loops controlling the carbon dynamics, could provide powerful tools to unravel this process interplay, to enable a direct comparison between model results and observations and to test alternative hypotheses concerning the causes and effects of extreme perturbations of the global carbon cycle and climate.

However, although state-of-the-art early diagenetic models are sophisticated and comprehensive enough to accurately reproduce observations and predict exchange fluxes (see e.g. Boudreau, 1997; Arndt et al., 2013), most paleoclimate models do not resolve the complexity of diagenetic processes. The primary constraint here is the high computation cost of simulating all of the essential redox and equilibrium reactions within marine sediments that control carbon burial and benthic recycling fluxes: a barrier that is exacerbated if a variety of benthic environments are to be spatially resolved. Instead, most models either neglect or only roughly approximate biogeochemical processes in the sediment and the benthic-pelagic coupling. In the following, we describe model approaches for shallow-water coral reefs and, similar to results of an earlier review of the coupling between benthic and pelagic biogeochemical models (Soetaert et al., 2000), four representations for deep-sea sediments characterised by different levels of complexity (Figs. 14 and 15).

### 3.3.1. Shallow-water carbonate sediments

Not all burial of carbonates takes place in the deep-sea; it is currently estimated that approximately an equal amount accumulates in shallow-water (neritic) environments (Milliman and Droxler, 1996; Vecsei, 2004). Neritic carbonates are primarily the product of seafloor dwelling calcifying organisms such as corals, echinoids, mollusks, benthic foraminifera, bivalves, sea urchins, and coralline algae, whose long-term accumulation can result in the formation of large carbonate banks or reefs (e.g. Perry et al., 2008). As today the surface ocean is largely oversaturated with respect to aragonite ( $\Omega_{Ar}$ ), most of the global shallow-water carbonate production (about 0.65–0.83 Pg  $\text{CaCO}_3 \text{ year}^{-1}$ , Vecsei (2004), i.e., 0.078–0.100 Pg C  $\text{year}^{-1}$ ) is retained and accumulates in coral-reef sediments. Additionally, particulate organic carbon flux and sedimentation rates are elevated in neritic environments therefore often leading to suboxic and anoxic conditions in the sediments. The current knowledge is that the rate of calcification is controlled by a combination of ambient factors such as aragonite saturation state of the seawater, temperature and light availability (e.g. Kleypas and Langdon, 2006; Tambutté et al., 2011; Jones et al., 2015). However, more research is needed in order to improve the understanding of the interplay of physico-chemical and biological factors controlling the formation and composition of shallow carbonates (Allemand et al., 2011). In some paleoclimate models, such as GENIE (Cui et al., 2013), MBM (Munhoven, 2007; Munhoven and François, 1996) and GEOCLIM (Donnadieu et al., 2006), shallow-water carbonate formation depends on the saturation state of the epicontinental ocean with respect to  $\text{CaCO}_3$ , as well as on the total shelf area available for the formation of carbonate platforms (Goddéris and Joachimski, 2004) and possibly the rate of sea-level change (Walker and Opdyke, 1995; Munhoven and François, 1996). The majority of paleoclimate models, however, ignore carbonate deposition in shallow-water environments because of resolution issues and the high computational requirements to model the involved suboxic and anoxic redox-reactions.

### 3.3.2. Deep-sea sediments

Paleoclimate models use a wide variety of approaches to represent ocean-sediment exchange fluxes. The most simple ones do not include any explicit sediment scheme, but simply assume that particulate fluxes reaching the bottom of the ocean degrade there, and the remineralisation or dissolution products return to the deepest model grid cells or boxes of the ocean. Remineralisation and/or dissolution may be either complete or partial. In the latter case, the non-degraded fraction is returned to the surface ocean, to mimic riverine input, thus avoiding the model drift because of global inventory changes. POC remineralisation rates may be dependent on oxygen concentrations in the grid cells just above the seafloor and PIC dissolution rates on the saturation state with respect to the saturation state of bottom waters. Some models (e.g. BICYCLE) use a restoration scheme, either based upon a prescribed history of the sedimentary lysocline, which is used as a proxy for the




Fig. 14. Schematic representation of the four different sediment approaches in paleoclimate models. Adapted from Soetaert et al. (2000). A: Reflective boundary; B: conservative/semi-reflective boundary; C: vertically-integrated, dynamic model; D: vertically resolved, diagenetic model (same as Fig. 12).

calcite saturation horizon (Köhler et al., 2005) or a reference deep-sea  $\text{CO}_3^{2-}$  concentration (Köhler et al., 2010). Even models with explicit representations of the surface sediments exhibit a large variety of configurations: along the vertical in the sediment column, complexity ranges from single-box surface mixed-layers (e.g., Munhoven and François, 1996) to well resolved sediment columns (e.g., 21 grid-points for the surface mixed-layer in MEDUSA, Munhoven, 2007). Underneath the mixed-layer, some models additionally track the history of preservation in synthetic sediment cores (e.g., Ridgwell and Hargreaves, 2007). The composition of the model sediments is also highly variable, encompassing the range from a minimalistic calcite-clay mixture (Munhoven and François, 1996; Zeebe, 2012) with an implicit, steady-state porewater  $[\text{CO}_3^{2-}]$  profile, to a composition that essentially reflects the ocean model tracer (e.g. DIC,  $\text{O}_2$ ,  $\text{PO}_4$ ) in porewaters and material fluxes (e.g. PIC, POC,  $\text{CaCO}_3$ , opal and clay) in the solid fraction (Bern 3D, Tschumi et al., 2011). Except for MEDUSA in MBM (Munhoven, 2007), no sediment model appears to explicitly consider aragonite as a sediment constituent. In other models, the entire aragonite flux is dissolved close to the sediment-water interface if bottom waters are undersaturated with respect to aragonite, while the flux is entirely

preserved in oversaturated bottom waters, possibly “converted” to calcite (e.g. in MBM, Munhoven and François, 1996). The various adopted approaches may be divided into four major classes, which we review in the following.

**3.3.2.1. Reflective boundary.** The Bern 2.5D model includes the sediment-water interface in form of a reflective boundary (Marchal et al., 1998b) and also the GENIE model provides this as an option (Ridgwell et al., 2007) (Fig. 14A). The deposition flux of PIC and POC that would settle onto the sediments is completely consumed in the deepest ocean cell, instantaneously releasing dissolved carbon and nutrients. This approach is, due to its computational efficiency, often used in global biogeochemical models (e.g. Najjar et al., 1992; Sarmiento et al., 1995; Yamanaka and Tajika, 1996). Usually the partitioning of the return flux (representing benthic transformations) is parameterised by using molar ratios for the particulates but could also be calculated based on steady-state diagenetic modelling (Soetaert and Herman, 1995; Regnier et al., 1997). As the reflective boundary approach does not model any benthic PIC and POC burial fluxes it overestimates benthic recycling fluxes and completely neglects the

<b>Conservative/semi-reflective boundary</b>			
e.g. in LOVECLIM (Goosse et al., 2010)			
$F_{S,bur} = F_{S,dep}$	( or $\alpha \cdot F_{S,dep}$ ) (1)		
$F_{D,rf} = 0$	( or $R \cdot (1 - \alpha) \cdot F_{S,dep}$ ) (2)		
$\alpha \in (0, 1]$ , (with $\alpha = 1$ for the conservative boundary)			
$C_{D,sed} = 0$	(3)		
<b>Reflective boundary</b>			
e.g., in Bern2.5D (Marchal et al., 1998b) and GENIE (Ridgwell et al., 2007):			
$F_{S,bur} = 0$	(4)		
$F_{D,rf} = g(F_{S,dep})$	( or $R \cdot F_{S,dep}$ ) (5)		
$C_{D,sed} = 0$	(6)		
<b>Vertically-integrated, dynamic model</b>			
e.g., in Bern3D (Tschumi et al., 2011), LOSCAR (Zeebe, 2012) or MBM1996 (Munhoven and François, 1996):			
$F_{S,bur} = F_{S,dep} - \sum_j Reac_j$	(7)		
$F_{D,rf} = \sum_j \beta_j Reac_j$	(8)		
$\frac{dC_{D,sed}(t)}{dt} = F_{D,in} - F_{D,out} + \sum_j (1 - \beta_j) Reac_j$	(9)		
<b>Vertically resolved, diagenetic model</b>			
General steady-state diagenetic equation for solid and dissolved species $C_i$ after Berner (1980). See e.g. GEOCLIM reloaded (Arndt et al., 2011):			
$\frac{\partial \xi C_i(t, z)}{\partial t} = 0 = -\frac{\partial}{\partial z} F + \xi \sum_j Reac_j$	(10)		
$= -\frac{\partial}{\partial z} \left( -\xi D_i \frac{\partial C_i}{\partial z} + \xi w C_i \right) + \xi \sum_j Reac_j$			
<b>Glossary</b>			
$S$	Solid species	$D$	Dissolved species
$F_{S,bur}$	Sediment burial flux of solids	$F_{i,dep}$	Bottom water deposition flux of $i$
$F_{D,rf}$	Dissolved return flux due to OM remineralisation	$C_{D,sed}$	Sediment concentration of D
$\alpha$	Fraction of solids preserved	$F_{D,in/out}$	General dissolved in/out-flow
$g()$	Steady state return flux	$R$	Stoichiometric ratio
$z$	Sediment depth	$\beta_j \in [0, 1]$	Dissolved fraction returned to ocean
$\sum_j Reac_j$	Sum of relevant production /consumption processes	$z_{max}$	Maximum sediment depth
$F$	Transport fluxes	$\xi$	Equals porosity $\phi$ for solutes and ( $1 - \phi$ ) for solids
$w$	advection rate	$D_i$	Diffusion coefficient

**Fig. 15.** Overview of applied model approaches to calculate burial of solid species (i.e. PIC and POC), return/recycling fluxes of dissolved species resulting from organic matter (OM) remineralisation and sediment concentrations of dissolved species.

strong coupling between POC and PIC fluxes through the effect of organic matter degradation on carbonate dissolution (Fig. 13). In addition, it does not account for the temporal storage of material in the sediment and the time delay between deposition and recycling flux. Therefore, this highly simplified approach cannot resolve the complex benthic-pelagic coupling.

**3.3.2.2. Conservative/semi-reflective boundary.** The conservative boundary approach (Fig. 14B) refers to biogeochemical models that impose burial fluxes and sediment-water exchange fluxes. In general, the burial flux of PIC and POC is set equal (or proportional) to their deposition flux. In addition, a no-flux boundary condition is applied for solute species, neglecting any exchange through the sediment-water interface. For instance LOVECLIM incorporates such a sediment model




Fig. 16. Model comparison of marine biogeochemistry with GENIE configurations using the Conservative and Reflective sediment approach for the modern and Eocene world. Modelled global oxygen (A) and phosphate (B) profiles. (C + D): Anomaly plots (Conservative minus Reflective) of bottom water oxygen concentration for modern (C) and Eocene (D) world.




Fig. 17. Model comparison of  $H_2S$  concentration during OAE2 with GENIE configurations using the conservative and reflective sediment approach. (A + B): Vertical profile of  $H_2S$  in the East Pacific Ocean ( $-90^\circ$  longitude). (C): Anomaly plots (Reflective minus Conservative) of photic zone euxinia (i.e.  $H_2S$  concentration for 80–200 m depth).

where a constant part of POC and PIC is preserved (Goosse et al., 2010). The conservative nature of this approach does not violate mass conservation and accounts for the retention capacity of sediments. Nevertheless, it neglects (or oversimplifies) the degradation of POC and the dissolution of PIC in marine sediments and thus overestimates (or crudely approximates) the burial fluxes. In addition, such a simplified approach does not represent the time-delayed recycling of nutrients and dissolved carbon and the impact of these fluxes on the biogeochemical

functioning of the ocean-atmosphere system. Another important caveat of this approach is that it cannot account for a change in speciation. Generally, the composition of the benthic return fluxes is fundamentally different from the composition of the deposition flux (e.g. Soetaert et al., 2000). In marine sediments, the coupled redox-reactions, mineral precipitation/dissolution or equilibrium reactions control the speciation. The exact composition of the total dissolved carbon flux, for instance, strongly depends on the vertical distribution of




Fig. 18. Idealised dependency of various sources of uncertainty on model complexity (i.e. components, resolution and represented processes). Blue arrows depict an improvement of input or parameter uncertainty using SA and empirical methods (see text) which results in a decrease in total model uncertainty (black arrows).

Source: Adapted from Solomatine and Wagener (2011).

biogeochemical reaction rates and their combined influence on the ambient pH. The conservative boundary approach does not capture this biogeochemical complexity and thus does not appropriately represent the sedimentary response.

In order to illustrate how different sediment boundary conditions affect biogeochemistry in the ocean we compare a GENIE set-up using the reflective boundary with the conservative boundary for two climate scenarios (see Box 2 for more information). The impact of including organic carbon burial on global mean water-column  $O_2$  and  $PO_4$  concentration during the Eocene is shown in Fig. 16 (A + B). Global deep water  $O_2$  concentration increases as POC reaching the seafloor is buried and not remineralised. In contrast, nutrient concentration in the deep ocean is decreasing as less  $PO_4$  is released to the ocean. But not just the global  $O_2$  concentration changes, also the spatial difference of bottom water oxygenation for the two sediment schemes varies significantly for the modern and the Eocene (Fig. 16C + D). Ocean redox differences are even more pronounced when applying the two sediment representations for the Late Cretaceous. Fig. 17 (A) highlights the problem of the reflective lower boundary by showing an unrealistically high concentration of  $H_2S$  at the seafloor. This result is an artifact of the lower boundary condition as all POC gets instantaneously remineralised at the seafloor. On the other hand, the conservative boundary (Fig. 17B) shows very little  $H_2S$  in the deeper ocean as it neglects completely the degradation of POC at the seafloor. However, the employed lower boundary condition does not only have implications on redox conditions at the bottom of the ocean but can also be seen in the photic zone (Fig. 17C).

**3.3.2.3. Vertically-integrated dynamic model.** In the vertically-integrated approach, the sediment is represented as a single box (Fig. 14C). The average concentration of the represented species in this box is calculated as the balance between the deposition and burial flux, as well as the sum of consumption processes. The diffusive flux of dissolved species through the sediment-water interface in turn equals the sum of consumption/production processes that are usually tightly linked to the transformation of particulate material (e.g. Maier-Reimer, 1993). The model thus neglects temporary storage of dissolved species and fluxes in porewaters. However, this approach is clearly superior to the two simpler approaches. It has the merit of simplicity and is computationally efficient. In addition, it also reproduces some of the complexity associated with the short- and long-term evolution of benthic recycling fluxes. Such an approach also allows differentiating between various fractions of organic matter (if POC is represented) and

therefore is able to resolve some of the biogeochemical complexity associated with the decrease of organic matter reactivity with sediment depth. Most paleo-EMICs incorporate a vertically-integrated sediment model for PIC only, sometimes considering oxic-only sediment respiration of organic carbon (compare Table 1). Bern 3D being a notable exception, as it includes a vertically-integrated dynamic model also considering oxic degradation and denitrification of organic carbon (Tschumi et al., 2011),

**3.3.2.4. Vertically resolved diagenetic model.** So-called diagenetic models provide the most robust description of the benthic-pelagic coupling (Fig. 14D). Those models solve the one-dimensional, fully coupled reaction-transport equation for solid and dissolved species (e.g. Berner, 1980b; Boudreau, 1997). This approach thus accounts for all important transport processes, such as burial, compaction, bioturbation, molecular diffusion and bioirrigation. In addition it resolves the fully coupled biogeochemical dynamics of the carbon, oxygen and nutrient cycles and the resulting characteristic redox-zonation of marine sediments (also compare Fig. 12). However, controversy still revolves around the formulations of organic matter degradation (e.g. Arndt et al., 2013) and calcite dissolution (e.g. Jourabchi et al., 2005). In addition, the parameterisation of diagenetic models requires a good understanding of diagenetic dynamics and careful consideration of the environmental conditions. For instance, rate constants that are typically used in state-of-the-art diagenetic models may predict the benthic response in the modern-day, well ventilated ocean, but might not be applicable under extreme environmental conditions such as OAEs and the PETM. The major drawback of those models is the computational cost associated with the computation of vertically-resolved reaction-transport equations for a number of interacting species. Therefore, paleoclimate models that include a diagenetic model generally reveal a very low spatial resolution of the benthic environment (e.g. GEOCLIM reloaded only resolves three sediment columns; Arndt et al., 2011) or use other methods to reduce computational demands: DCESS for instance uses a semi-analytical, iterative approach considering  $CaCO_3$  dissolution and (oxic and anoxic) organic matter remineralisation (Shaffer et al., 2008). One exception here is the early diagenesis model MEDUSA which is coupled to the multi-box model MBM (Munhoven, 2007). MEDUSA operates in a fully transient way at 100 m depth intervals over the whole model sea-floor, in five regions, totalling 304 columns.

### 3.3.3. Conclusion

Marine sediments represent the largest reservoir of carbon among the exogenic reservoirs (Mackenzie et al., 2004). The assessment of the response of the ocean to variabilities in atmospheric  $CO_2$  concentrations requires a robust quantification of the benthic-pelagic coupling and the sedimentary carbon sink (Archer and Maier-Reimer, 1994; Archer et al., 1998; Sigman et al., 1998; Heinze et al., 1999). However, it appears that convenience rather than a careful mechanistic representation and the ability of the approach to provide an answer to the problem guides the choice of the lower boundary condition for the ocean model. Paleoclimate modelling has developed to a stage where increasingly complex and multi-dimensional ocean, atmosphere and continental vegetation models are coupled (e.g. McGuffie and Henderson-Sellers, 2005; Randall et al., 2007). Yet, compared to these developments, considerably less effort has been devoted to the coupling between ocean and sediment models. However, sophisticated, comprehensive and carefully calibrated and tested diagenetic models (e.g. Aguilera et al., 2005; Archer et al., 1998; Cappellen and Wang, 1996; Soetaert et al., 1996), as well as computationally efficient pseudo dynamic approaches (e.g. Ruardij and Van Raaphorst, 1995; Arndt and Regnier, 2007) are now available and could be incorporated into paleoclimate models in numerically efficient ways, such as for instance look-up tables (see e.g. Ridgwell and Hargreaves, 2007) and neural networks (see Section 4.3). Ultimately, our ability to understand past climate change critically

depends on a better quantification of the sedimentary carbon sink and its response to extreme environmental conditions (Berner et al., 1989; Archer and Maier-Reimer, 1994; Mackenzie et al., 2004; Ridgwell and Zeebe, 2005).

#### 4. Discussion and conclusions

Marine carbon cycling involves complex interactions between biology, chemistry, and physics and represents a dynamic system that exhibits feedbacks with changes in climate. The mechanistic understanding of processes involved continues to improve and an increasing number of paleoclimate models incorporating marine carbon cycle dynamics are becoming available. However, the organic and inorganic carbon cycling in the ocean and the benthic-pelagic coupling are still widely represented by highly simplified approaches and are therefore of limited transferability across time and space. Progress in understanding past climate variations will crucially depend on the combined use of different representations (e.g. conceptual and mechanistic) of the surficial carbon cycle and the quantification of related model uncertainties.

In this final section we highlight the importance of using models of different complexities (Section 4.1), provide suggestions how these models can be applied to explore and quantify different model uncertainties (Section 4.2), and identify two major challenges to help direct future research for the paleoclimate (modelling) community (Section 4.3).

##### 4.1. The importance of different models

Fundamentally, numerical models are always approximations of the complex, real Earth system and all assumptions are erroneous on some level (Oreskes et al., 1994). For instance, assuming a reflective sediment-water interface is appropriate when investigating carbon cycle processes on time scales shorter than 1000 years but is misleading when studying longer time scales. A box model approach is helpful when trying to isolate the dominant process in an observed global or large-scale output but not very helpful if one is interested in a more detailed (spatial) analysis of a problem (e.g. modelling marine ecology which is dependent on local transport and mixing processes and the spatial resolution of the ocean) or contrasting with regionally-specific observations.

Both model types (i.e. structurally simple or conceptual models and more coupled or mechanistic models) have their advantages and disadvantages (compare e.g. Nihoul, 1994). The structural simplicity of box models considerably reduces the models dependency on initial and boundary conditions and the model is easier to constrain as it includes fewer parameters and variables. Due to their lower computational demand box models can be used for large-ensemble experiments needed to address important questions regarding uncertainty quantification (see Section 4.2). Also, the output is less complex, easier to interpret and therefore may provide a clearer understanding of the dominant process. However, there is a higher possibility of misinterpreting the *real* process if it is actually the product of several interacting processes not represented in the model. Furthermore, the simplicity of the model (in terms of resolution and represented processes) usually restricts the development of emergent behaviours and in the case of ocean circulation and prescribed water mass exchanges between boxes, there can be no emergent behaviour (in circulation). More coupled or mechanistic models provide a slightly more complete view of the real system and have a greater degree of freedom in their dynamical properties. However, simply including more and more complexity (in the sense of additional mechanisms) does not guarantee an improvement of the predictive ability of the model. It may even reduce it, if the new representation is based on over fitting imprecisely known free parameters to limited observations (Davies, 1994). Thus, a crucial step is to show that the representation of the new mechanism has an acceptable

level of accuracy over a range of conditions.

Improving mechanistic parameterisations of key processes is one of the main challenges that hinder better understanding in Earth sciences. That does not, however, undermine the value of non-mechanistic models (e.g. conceptual, mathematical, statistical or numerical). Starting an investigation with a simple model and gradually increasing its complexity can reveal emerging model and system behaviours which might have been overlooked when employing the most complex model alone (compare Marinov et al., 2008, for an example). Also, how processes are represented even within the same model structure can vary and testing different assumptions can lead to improved understanding of the system, as has been demonstrated in this review. Hence, one way forward is rather than solely using existing, static numerical representations for the biological pump and trying to reproduce certain paleo-observations as perfectly as possible (which is in essence often just a fitting exercise), paleoclimate models can be used to explore alternative methodologies and biogeochemical mechanisms and hence map out the range of dynamical behaviour of the biological carbon pump that is possible. While not then necessarily fitting the observations of any one particular geologic event, assessing the behaviours of different incarnations (of the same model) may provide insight into other events or potentially, future responses (for which there is no data constraints). The resulting range of states or responses to perturbation, can also provide insight into model (structural) uncertainty, which we discuss next.

##### 4.2. Quantifying uncertainty

Beside being ideal tools for testing our understanding of the biological carbon pump and for exploring the long-term carbon cycle evolution, paleoclimate models should further be used to investigate uncertainties related to modelling climate and marine carbon cycle feedbacks and to identify which processes have the greatest influence upon model predictions. Currently uncertainty estimates for the climate-carbon cycle response are primarily done using different future emission scenarios (e.g. Plattner et al., 2008; Eby et al., 2009; Zickfeld et al., 2013) or, where possible, by comparing model results with observational uncertainty estimates (e.g. Najjar et al., 2007; Eby et al., 2013). In the case of model intercomparison projects mainly simplified characteristics such as the ensemble standard deviation and range are used (e.g. Najjar et al., 2007; Plattner et al., 2008; Zickfeld et al., 2013) or EMIC results are compared with results obtained from GCMs (e.g. Petoukhov et al., 2005; Friedlingstein et al., 2006; Plattner et al., 2008).

Uncertainty in (Earth system) modelling can generally be considered as a lack of knowledge or information concerning the processes involved and comes from a variety of sources (Solomatine and Wagener, 2011). First, there is input uncertainty, that is the uncertainty caused by errors in the boundary conditions, such as continental configuration, bathymetry, oceanic nutrient concentration and atmospheric CO<sub>2</sub> forcing. Second, there is parameter uncertainty, introduced through uncertain estimates of model parameters or because the optimal parameter set is ambiguous. And third, model structural uncertainty, resulting from simplifications, discretisations, inadequacies or ambiguities in the numerical representation of the real process. The different sources of uncertainty generally vary with model complexity (Fig. 18). Ideally, as more processes are described in the model (i.e. an increase in model complexity) the structural uncertainty of the model decreases. However, at the same time more parameters and inputs are needed to describe and/or constrain these processes, therefore increasing input and parameter uncertainty. Due to this trade-off between different complexities there is theoretically an optimal model for every given real world problem characterised by minimal total uncertainty. Obviously, this is an idealised example and in reality it is not possible to decide which model is the optimal one, especially in the case of paleoclimate modelling where validation against observations is limited.

One strategy to explore and quantify model uncertainty that can

always be applied is sensitivity analysis (SA). Pianosi et al. (2014) deliver a comprehensive review of SA methods most widely used in other environmental modelling fields. They also provide practical guidelines for choosing the most appropriate SA method for a specific problem, depending on the purpose of the analysis and the computational complexity of the method. Sensitivity analysis in combination with empirical approaches can also be used to iteratively reduce the uncertainty for a given numerical model (compare e.g. Bradley et al., 2016). For instance in the case of parameter uncertainty, SA can identify which parameters have the greatest influence upon model output and how constrained their values are. Mechanistic parameters with explicit relevance to biology or physics (such as degradation rate constants, activation energies and temperature dependencies) may then be better constrained empirically using laboratory experiments or more observations, thus decreasing parameter uncertainty (compare Fig. 18). However, more conceptual parameters (e.g. Martin's b-value), that implicitly account for various processes, are more specific to individual model formulations and have to be determined by fitting each model to observations. The lack of correlation between these abstract parameters and real biogeochemical processes leads to an increase in parameter uncertainty (i.e. decrease in predictive ability) if the numerical model is applied under different environmental conditions. SA should be applied here to quantify this uncertainty.

Currently most studies treat only parameter uncertainty as source of uncertainty. However, research in hydrological modelling has shown that input or structural uncertainty can be even more important (Wagener and Gupta, 2005; Solomatine and Wagener, 2011). Therefore, other questions concerning input and structural uncertainty should be explored more often using SA: How much does it matter if we change boundary conditions or forcings? What is the uncertainty in model results due to missing or oversimplified biogeochemistry? The computational efficiency of paleoclimate models (especially of box models) facilitates large simulation ensembles and systematic sensitivity studies thus allowing the estimation of these uncertainties using objective statistical methods. Kriest et al. (2010) and Kriest et al. (2012) for example present extensive sensitivity experiments to assess the skill of different global marine biogeochemical models for the modern ocean.

#### 4.3. Outstanding modelling issues

As shown in this study, a major problem is the lack of the existence of a fundamental theoretical framework for describing the particulate organic carbon flux within the water column, and hence the limited ability to directly derive a parameterisation which is applicable under changing environmental conditions (A universal single representation would not be possible anyway, as illustrated for instance by physical ocean circulation modelling where although the fundamental fluid dynamics on a rotating sphere is known, practical constraints of model resolution and time-step lead to different approximations and parameterisations.). Different generic algorithms for this parameterisation, based on current process understanding, are needed to resolve changes in the efficiency of the biological pump which is observed over time (e.g., Sigman and Boyle, 2000; Kohfeld et al., 2005; John et al., 2014)

## Appendix A. Earth system model applications

Table 5

Paleoclimate model applications (EMICs) (ordered from higher to lower ocean resolution) with a relation to the biological pump. Study-time reflects the broad modelled time in the experiments (not the official timing of the event).

Model	Event	Study-time	References
<b>EMICs</b>			
Bern 2.5D	Younger Dryas Heinrich events	ca. 13.0–11.0 kyr BP ca. 60–12 kyr BP	Marchal et al. (1999a,b, 2001) Marchal et al. (1998a)

and in space (e.g., Weber et al., 2016). Although the basic power-law or double-exponential decrease in POC flux with depth has been widely used and characterised, there is a strong need to link changes in downward particulate flux to the mechanistic understanding of the underlying processes. The results of different representations should be compared against each other as discussed above.

Another outstanding issue are the current oversimplified sediment representations in paleoclimate models, which are often not able to model the properties needed for a comparison to proxy-observations. To overcome comparable problems, offline coupling approaches are often used. For instance, sophisticated atmospheric models such as FOAM are often run for a wide range of different atmospheric CO<sub>2</sub> concentrations. Based on these stand-alone simulations, look-up tables are created that link the modelled climatic conditions (e.g. precipitation, temperature) to atmospheric CO<sub>2</sub> (e.g., Donnadieu et al., 2006). These look-up tables are numerically efficient representations of the full model that can subsequently be used in EMIC simulations to simulate the climatic response to changing atmospheric CO<sub>2</sub> concentrations. Artificial neural networks (which represent more sophisticated look-up tables or transfer functions that allow linking a number of input parameters to a number of output parameters) can be used to encapsulate a numerically cost intensive model in a similar way. A neural network encapsulation of sediment dynamics has to be trained using a large set of simulated bottom water conditions and related burial and benthic recycling fluxes calculated by a full diagenetic model. Instead of following static, human prescribed rules, neural networks have the ability to learn automatically the underlying relationship from the training set and are practically able to approximate any relationship between input and output properties (Melesse and Hanley, 2005; Bernhardt, 2008). These approaches thus allow a full, dynamic coupling of both models at a significantly reduced computational cost. While a neural network in effect becomes a 'black box' and loses the ability to be able to ascribe causes to consequences, it should be noted that the existence of the underlying mechanistic model on which a neural network has been trained allows for the dynamics of the system as a whole to still be picked apart/understood; i.e. unlike training a neural network on data when nothing per se can then be learned about processes, the underlying mechanistic model can be employed alongside (or in limited cases, in lieu of) the neural network to ensure that a mechanistic interpretation can always be made.

#### Acknowledgements

We would like to thank two anonymous reviewers for their constructive critiques and suggestions that have greatly improved this paper. We are also grateful to Francesca Pianosi and Thorsten Wagener for helpful insights into quantifying uncertainty and to Sarah Greene for stimulating discussions. DH is supported by a graduate teaching studentship by the University of Bristol. SA is supported by funding from the European Unions Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 643052 (C-CASCADES). JDW and AR acknowledge funding from the EU grant ERC-2013-CoG-617313. GM is a Research Associate with the Belgian Fonds de la Recherche Scientifique-FNRS.

Bern 3D	& last deglaciation		
	Holocene	ca. 8 kyr BP–present	Menviel and Joos (2012), Brovkin et al. (2016)
	LGM and deglaciation	20.0 kyr BP–present	Tschumi et al. (2008, 2011 )
	Last glacial period	ca. 70–30 kyr BP	Parekh et al. (2008)
CLIMBER-2	Last glacial cycle	125 kyr BP–present	Menviel et al. (2012)
	Emian interglacial	126–115 kyr BP	Brovkin et al. (2016)
	Holocene	ca. 8 kyr BP–present	Kleinen et al. (2015), Brovkin et al. (2016)
	8.2 kyr BP event	8.2 kyr BP	Brovkin et al. (2002a)
GENIE	Last deglaciation	ca. 18–9 kyr BP	Bouttes et al. (2012a)
	LGM	ca. 21 kyr BP	Brovkin et al. (2002b, 2007 ), Bouttes et al. (2009)
	Emian interglacial	ca. 125 kyr BP	Bouttes et al. (2010, 2011, 2012b)
	Glacial cycles	ca. 126–116 kyr BP	Duplessy et al. (2007)
	MIS 11 interglacial	126 kyr BP–present	Kleinen et al. (2015), Brovkin et al. (2016)
	Holocene	420–380 kyr BP	Brovkin et al. (2012)
	Younger Dryas	8–0.5 kyr BP	Kleinen et al. (2015)
	Emian interglacial	13.5–11.5 kyr BP	Brovkin et al. (2016)
	Eocene hyperthermal	126–115 kyr BP	Singarayer et al. (2008)
	Early Eocene	49.2 Ma	Brovkin et al. (2016)
GEOCLIM	PETM	ca. 50 Ma	Kirtland Turner and Ridgwell (2013)
	OAE2	ca. 55 Ma	Norris et al. (2013)
	Cretaceous	ca. 55.8 Ma	Meyer and Kump (2008), Pälike et al. (2012)
	End-Permian		Anagnostou et al. (2016)
	Last 400 Ma	ca. 55.8 Ma	Ridgwell (2007), Panchuk et al. (2008)
	Cretaceous	400 Ma–present	Ridgwell and Schmidt (2010), Cui et al. (2011)
	Jurassic	145.5–65.5 Ma	Kirtland Turner and Ridgwell (2016)
	Triassic	199.6–145.5 Ma	Zeebe et al. (2016)
	Late Neoproterozoic	251.0–199.6 Ma	Monteiro et al. (2012)
	Neoproterozoic	580.0 Ma	Meyer and Kump (2008)
LOVECLIM	Last deglaciation	1000–542.0 Ma	Meyer and Kump (2008), Meyer et al. (2008)
	LGM	ca. 21–10 kyr BP	Cui et al. (, 2013, 2015 )
	Interglacials	ca. 22.5 kyr BP	Goodwin et al. (2009)
	Emian interglacial	Universally	Donnadieu et al. (2006)
MESMO	Younger Dryas	ca. 129–118 kyr BP	Donnadieu et al. (2006)
	Heinrich 1	ca. 12.9–11.7 kyr BP	Donnadieu et al. (2006), Godd�ris et al. (2008)
	LGM	ca. 17.5–14.5 kyr BP	Donnadieu et al. (2004)
	Glacial cycles	ca. 22.5 kyr BP	Le Hir et al. (2008)
MoBidiC	LGM	ca. 400 kry BP–present	Menviel et al. (2011)
	Heinrich events	ca. 22.5 kyr BP	Menviel et al. (2008a,b, 2014)
	& Last deglaciation	ca. 60–12 kyr BP	Menviel et al. (2010)
	UVic	ca. 20–15 kyr BP	Duplessy et al. (2007)
UVic	Last deglaciation	23 kyr BP–present	Matsumoto and Yokoyama (2013)
	LGM	ca. 21 kyr BP	Matsumoto and Yokoyama (2013)
	DO-event 8 & 12	ca. 37 & 45 kyr BP	Sun and Matsumoto (2010), Matsumoto et al. (2014)
	DO-event	Universally	Ushie and Matsumoto (2012)
	Panama seaway closure	ca. 14–3 Ma	Crucifix (2005)
	Antarctic Glaciation	ca. 33.7 Ma	Crucifix (2005)
	PETM	ca. 55.8 Ma	Schmittner and Lund (2015)
	End-Permian	ca. 252 Ma	Simmons et al. (2015)
			Fraille et al. (2009), Menviel et al. (2014)
			Schmittner and Somes (2016)
		Ewen et al. (2004)	
		Schmittner (2005), Schmittner et al. (2007)	
		Schneider and Schmittner (2006)	
		Pagani et al. (2011)	
		Meissner et al. (2014), Bralower et al. (2014)	
		Alexander et al. (2015)	
		Montenegro et al. (2011)	

Table 6

Paleoclimate model applications (box models) (ordered from higher to lower ocean resolution) with a relation to the biological pump. Study-time reflects the broad modelled time in the experiments (not the official timing of the event).

Model	Event	Study-time	References
<b>Box models</b>			
BICYCLE	Last deglaciation	21–10 kyr BP	Köhler et al. (2005)
	Glacial cycles	120 kyr BP–present	Köhler et al. (2006)
	Glacial cycles	740 kyr BP–present	Köhler and Fischer (2006), Köhler et al. (2010)
	Mid Pleistocene Transition	2 Ma–present	Köhler and Bintanja (2008)
CYCLOPS	LGM	ca. 22.5 kyr BP	Keir (1988), Michel et al. (1995), Sigman et al. (2003) Galbraith et al. (2015)
	Last deglaciation	ca. 30 kyr BP–present	Keir (1991), Sigman et al. (1998)
DCESS	Glacial cycle	150 kyr BP–present	Keir (1990), Hain et al. (2010)
	LGM	ca. 22.5 kyr BP	Lambert et al. (2015)
	PETM	ca. 55.8 Ma	Shaffer et al. (2016)
LOSCAR	Mid-Miocene	16.8–13.8 Ma	Armstrong McKay et al. (2014)
	MECO	ca. 40 Ma	Sluijs et al. (2013)
	Paleocene-Eocene	62–48 Ma	Komar et al. (2013)
	Early Eocene	ca. 55 Ma	Pälike et al. (2012)
	PETM	ca. 55.8 Ma	Zeebe and Zachos (2007), Zeebe et al. (2009) Uchikawa and Zeebe (2010), Komar and Zeebe (2011) Zeebe and Zachos (2013), Penman et al. (2014) Zeebe et al. (2016)
MBM	Glacial cycle	ca. 150 kyr BP–present	Munhoven and François (1994, 1996) Munhoven (2007)
PANDORA	Glacial-interglacial CO <sub>2</sub>	Universally	Broecker and Peng (1986, 1987, 1989) Broecker et al. (1990)
SUE	Since LGM	ca. 22.5 kyr BP–today	Ridgwell et al. (2003b)
	Glacial cycles	400 kyr BP–present	Ridgwell (2001), Ridgwell et al. (2002)
	Snowball Earth	Around 650 Ma	Ridgwell et al. (2003a)
	Phanerozoic	600 Ma–present	Ridgwell (2005)

## References

- Aguilera, D.R.D.R., Jourabchi, P.P., Spiteri, C.C., Regnier, P.P., 2005. A knowledge-based reactive transport approach for the simulation of biogeochemical dynamics in earth systems. *Geochem. Geophys. Geosyst.* 6 <http://dx.doi.org/10.1029/2004GC000899>. <http://onlinelibrary.wiley.com/doi/10.1029/2004GC000899/abstract> Q07012.
- Alexander, K.K., Meissner, K.J.K.J., Bralower, T.J.T.J., 2015. Sudden spreading of corrosive bottom water during the Palaeocene-Eocene Thermal Maximum. *Nat. Geosci.* 8, 458–461. <http://dx.doi.org/10.1038/ngeo2430>. <http://www.nature.com/ngeo/journal/v8/n6/abs/ngeo2430.html> 00001.
- Allredge, A.L.A.L., Cohen, Y.Y., 1987. Can microscale chemical patches persist in the sea? Microelectrode study of marine snow, fecal pellets. *Science* 235, 689–691. <http://dx.doi.org/10.1126/science.235.4789.689>. <http://www.sciencemag.org/content/235/4789/689>.
- Allemand, D.D., Tambutté, E.E., Zoccola, D.D., Tambutté, S.S., 2011. Coral calcification, cells to reefs. In: Dubinsky, Z.Z., Stambler, N.N. (Eds.), *Coral Reefs: An Ecosystem in Transition*. Springer Netherlands, pp. 119–150. [http://link.springer.com/chapter/10.1007/978-94-007-0114-4\\_9](http://link.springer.com/chapter/10.1007/978-94-007-0114-4_9).
- Alverson, K.D.K.D., Bradley, R.S.R.S., Pedersen, T.F.T.F., 2003. *Paleoclimate, Global Change and the Future*. Springer Science & Business Media.
- Anagnostou, E.E., John, E.H.E.H., Edgar, K.M.K.M., Foster, G.L.G.L., Ridgwell, A.A., Inglis, G.N.G.N., Pancost, R.D.R.D., Lunt, D.J.D.J., Pearson, P.N.P.N., 2016. Changing atmospheric CO<sub>2</sub> concentration was the primary driver of early Cenozoic climate. *Nature*. <http://dx.doi.org/10.1038/nature17423>. <http://www.nature.com/nature/journal/vaop/ncurrent/full/nature17423.html> advance online publication 00000.
- Anderson, L.A.L.A., Sarmiento, J.L.J.L., 1994. Redfield ratios of remineralization determined by nutrient data analysis. *Global Biogeochem. Cycles* 8, 65–80. <http://dx.doi.org/10.1029/93GB03318>. <http://onlinelibrary.wiley.com/doi/10.1029/93GB03318/abstract>.
- Andersson, J.H.J.H., Wijsman, J.W.M.J.W.M., Herman, P.M.J.P.M.J., Middelburg, J.J.J.J., Soetaert, K.K., Heip, C.C., 2004. Respiration patterns in the deep ocean. *Geophys. Res. Lett.* 31 <http://dx.doi.org/10.1029/2003GL018756>. <http://onlinelibrary.wiley.com/doi/10.1029/2003GL018756/abstract>.
- Archer, D.D., 1996a. A data-driven model of the global calcite lysocline. *Global Biogeochem. Cycles* 10, 511–526. <http://dx.doi.org/10.1029/96GB01521>. <http://onlinelibrary.wiley.com/doi/10.1029/96GB01521/abstract>.
- Archer, D.D., Khesghi, H.H., Maier-Reimer, E.E., 1998. Dynamics of fossil fuel CO<sub>2</sub> neutralization by marine CaCO<sub>3</sub>. *Global Biogeochem. Cycles* 12, 259–276. <http://dx.doi.org/10.1029/98GB00744>. <http://onlinelibrary.wiley.com/doi/10.1029/98GB00744/abstract>.
- Archer, D.D., Maier-Reimer, E.E., 1994. Effect of deep-sea sedimentary calcite preservation on atmospheric CO<sub>2</sub> concentration. *Nature* 367, 260–263. <http://dx.doi.org/10.1038/367260a0>.
- Archer, D.D., Winguth, A.A., Lea, D.D., Mahowald, N.N., 2000a. What caused the glacial/interglacial atmospheric pCO<sub>2</sub> cycles? *Rev. Geophys.* 38, 159–189. <http://dx.doi.org/10.1029/1999RG000066>. <http://onlinelibrary.wiley.com/doi/10.1029/1999RG000066/abstract>.
- Archer, D.E.D.E., 1991. Modeling the calcite lysocline. *J. Geophys. Res.* 96, 17037–17050.
- Archer, D.E.D.E., 1996b. An atlas of the distribution of calcium carbonate in sediments of the deep sea. *Global Biogeochem. Cycles* 10, 159–174.
- Archer, D.E.D.E., Eshel, G.G., Winguth, A.A., Broecker, W.W., Pierrehumbert, R.R., Tobis, M.M., Jacob, R.R., 2000b. Atmospheric pCO<sub>2</sub> sensitivity to the biological pump in the ocean. *Global Biogeochem. Cy.* 14, 1219–1230. <http://dx.doi.org/10.1029/1999GB001216>.
- Armstrong, R.A.R.A., Lee, C.C., Hedges, J.J.J.J., Honjo, S.S., Wakeham, S.G.S.G., 2001. A new, mechanistic model for organic carbon fluxes in the ocean based on the quantitative association of POC with ballast minerals. *Deep-Sea Res. II Top. Stud. Oceanogr.* 49, 219–236. [http://dx.doi.org/10.1016/S0967-0645\(01\)00101-1](http://dx.doi.org/10.1016/S0967-0645(01)00101-1). <http://www.sciencedirect.com/science/article/pii/S0967064501001011>.
- Armstrong McKay, D.I.D.I., Tyrrell, T.T., Wilson, P.A.P.A., Foster, G.L.G.L., 2014. Estimating the impact of the cryptic degassing of Large Igneous Provinces: a mid-Miocene case-study. *Earth Planet. Sci. Lett.* 403, 254–262. <http://dx.doi.org/10.1016/j.epsl.2014.06.040>. <http://www.sciencedirect.com/science/article/pii/S0012821X14004257>.
- Arndt, S.S., Hetzel, A.A., Brumsack, H.-J.H.-J., 2009. Evolution of organic matter degradation in Cretaceous black shales inferred from authigenic barite: a reaction-transport model. *Geochim. Cosmochim. Acta* 73, 2000–2022. <http://dx.doi.org/10.1016/j.gca.2009.01.018>. <http://www.sciencedirect.com/science/article/pii/S0016703709000477>.
- Arndt, S.S., Jørgensen, B.B., LaRowe, D.D., Middelburg, J.J., Pancost, R.R., Regnier, P.P., 2013. Quantifying the degradation of organic matter in marine sediments: a review and synthesis. *Earth Sci. Rev.* 123, 53–86. <http://dx.doi.org/10.1016/j.earscirev.2013.02.008>. <http://www.sciencedirect.com/science/article/pii/S0012825213000512>.
- Arndt, S.S., Regnier, P.P., 2007. A model for the benthic-pelagic coupling of silica in estuarine ecosystems: sensitivity analysis and system scale simulation. *Biogeochem. Discuss.* 4, 747–796. <http://hal.archives-ouvertes.fr/hal-00297877>.
- Arndt, S.S., Regnier, P.P., Goddérès, Y.Y., Donnadiéu, Y.Y., 2011. GEOCLIM reloaded (v 1.0): a new coupled earth system model for past climate change. *Geosci. Model Dev.*

- 4, 451–481. <http://dx.doi.org/10.5194/gmd-4-451-2011>. <http://www.geosci-model-dev.net/4/451/2011/>.
- Arrieta, J.M.J.M., Mayol, E.E., Hansman, R.L.R.L., Herndl, G.J.G.J., Dittmar, T.T., Duarte, C.M.C.M., 2015. Dilution limits dissolved organic carbon utilization in the deep ocean. *Science* 348, 331–333. <http://dx.doi.org/10.1126/science.1258955>. <http://www.sciencemag.org/content/348/6232/331>.
- Arthur, M.A.M.A., Dean, W.W., Pollastro, R.R., Scholle, P.P., Claypool, G.G., et al., 1985. Comparative geochemical and mineralogical studies of two cyclic transgressive pelagic limestone units, Cretaceous Western Interior Basin, U.S. In: Pratt, L.M.L.M. (Ed.), *Fine-Grained Deposits and Biofacies of the Cretaceous Western Interior Seaway: Evidence of Cyclic Sedimentary Processes*. Soc. For Sediment. Geol. pp. 16–27. [http://archives.datapages.com/data/sepm\\_sp/fg4/Comparative\\_Geochemical.htm](http://archives.datapages.com/data/sepm_sp/fg4/Comparative_Geochemical.htm).
- Arthur, M.A.M.A., Dean, W.E.W.E., Pratt, L.M.L.M., 1988. Geochemical and climatic effects of increased marine organic carbon burial at the Cenomanian/Turonian boundary. *Nature* 335, 714–717. <http://dx.doi.org/10.1038/335714a0>. <http://www.nature.com/nature/journal/v335/n6192/abs/335714a0.html>.
- Arvidson, R.S.R.S., Mackenzie, F.T.F.T., Berner, R.A.R.A., 2014. The sensitivity of the Phanerozoic inorganic carbon system to the onset of pelagic sedimentation. *Aquat. Geochem.* 20, 343–362. <http://dx.doi.org/10.1007/s10498-013-9224-5>.
- Arvidson, R.S.R.S., Mackenzie, F.T.F.T., Guidry, M.M., 2006. MAGic: a Phanerozoic model for the geochemical cycling of major rock-forming components. *Am. J. Sci.* 306, 135–190. <http://dx.doi.org/10.2475/ajs.306.3.135>.
- Bacastow, R.B.R.B., Maier-Reimer, E.E., 1990. Ocean-circulation model of the carbon cycle. *Clim. Dynam.* 4, 95–125. <http://dx.doi.org/10.1007/BF00208905>.
- Barker, S.S., Higgins, J.A.J.A., Elderfield, H.H., 2003. The future of the carbon cycle: review, calcification response, ballast and feedback on atmospheric CO<sub>2</sub>. *Philos. Trans. R. Soc. Lond. A Math. Phys. Eng. Sci.* 361, 1977–1999. <http://dx.doi.org/10.1098/rsta.2003.1238>. <http://rsta.royalsocietypublishing.org/content/361/1810/1977>.
- Berelson, W.M.W.M., 2001. Particle settling rates increase with depth in the ocean. *Deep-Sea Res. II Top. Stud. Oceanogr.* 49, 237–251. [http://dx.doi.org/10.1016/S0967-0645\(01\)00102-3](http://dx.doi.org/10.1016/S0967-0645(01)00102-3). <http://www.sciencedirect.com/science/article/pii/S0967064501001023>.
- Berger, W.H.W.H., 1967. Foraminiferal ooze: solution at depths. *Science* 156, 383–385. <http://dx.doi.org/10.1126/science.156.3773.383>. <http://www.sciencemag.org/content/156/3773/383>.
- Berger, W.H.W.H., 1982. Increase of carbon dioxide in the atmosphere during deglaciation: the coral reef hypothesis. *Naturwissenschaften* 69, 87–88. <http://dx.doi.org/10.1007/BF00441228>.
- Berner, R.R., 1977. *Sedimentation and Dissolution of Pteropods in the Ocean, the Fate of Fossil Fuel CO<sub>2</sub> in the Oceans*. Plenum Press, New York, pp. 243–260.
- Berner, R.A.R.A., 1964. An idealized model of dissolved sulfate distribution in recent sediments. *Geochim. Cosmochim. Acta* 28, 1497–1503.
- Berner, R.A.R.A., 1980b. *Early Diagenesis: A Theoretical Approach*. Princeton University Press.
- Berner, R.A.R.A., 1980a. A rate model for organic matter decomposition during bacterial sulfate reduction in marine sediments. In: *Biogéochimie De La Matière Organique à L'interface Eau-Sédiment Marin*. Colloques Internationaux du CNRS. vol. 293. Centre National de la Recherche Scientifique, Paris, pp. 35–44.
- Berner, R.A.R.A., 1990. Atmospheric carbon dioxide levels over Phanerozoic time. *Science* 249, 1382–1386.
- Berner, R.A.R.A., 1991. A model for atmospheric CO<sub>2</sub> over Phanerozoic time. *Am. J. Sci.* 291, 339–376. <http://dx.doi.org/10.2475/ajs.291.4.339>.
- Berner, R.A.R.A., 1994. GEOCARB II: a revised model of atmospheric CO<sub>2</sub> over Phanerozoic time. *Am. J. Sci.* 294, 56–91. <http://dx.doi.org/10.2475/ajs.294.1.56>.
- Berner, R.A.R.A., Caldeira, K.K., 1997. The need for mass balance and feedback in the geochemical carbon cycle. *Geology* 25, 955–956. [http://dx.doi.org/10.1130/0091-7613\(1997\)025<0955:TNFMBIA>2.3.CO;2](http://dx.doi.org/10.1130/0091-7613(1997)025<0955:TNFMBIA>2.3.CO;2).
- Berner, R.A.R.A., Canfield, D.E.D.E., et al., 1989. A new model for atmospheric oxygen over Phanerozoic time. *Am. J. Sci.* 289, 333–361.
- Berner, R.A.R.A., Honjo, S.S., 1981. Pelagic sedimentation of aragonite: its geochemical significance. *Science* 211, 940–942. <http://dx.doi.org/10.1126/science.211.4485.940>. <http://www.sciencemag.org/content/211/4485/940>.
- Berner, R.A.R.A., Kothavala, Z.Z., 2001. GEOCARB III: a revised model of atmospheric CO<sub>2</sub> over Phanerozoic time. *Am. J. Sci.* 301, 182–204. <http://dx.doi.org/10.2475/ajs.301.2.182>.
- Berner, R.A.R.A., Lasaga, A.C.A.C., Garrels, R.M.R.M., 1983. The carbonate-silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years. *Am. J. Sci.* 283, 641–683. <http://dx.doi.org/10.2475/ajs.283.7.641>.
- Berner, R.A.R.A., Morse, J.W.J.W., 1974. Dissolution kinetics of calcium carbonate in sea water; IV, theory of calcite dissolution. *Am. J. Sci.* 274, 108–134. <http://dx.doi.org/10.2475/ajs.274.2.108>. <http://www.ajsonline.org/content/274/2/108>.
- Bernhardt, K.K., 2008. Finding alternatives and reduced formulations for process-based models. *Evol. Comput.* 16, 63–88. <http://dx.doi.org/10.1162/evco.2008.16.1.63>. 00008.
- Biddanda, B.B., Benner, R.R., 1997. Carbon, nitrogen, and carbohydrate fluxes during the production of particulate and dissolved organic matter by marine phytoplankton. *Limnol. Oceanogr.* 42, 506–518.
- Bishop, J.J., 1989. Regional extremes in particulate matter composition and flux: effects on the chemistry of the ocean interior. In: *Productivity of the Ocean: Present and Past*, 44, pp. 117–137.
- Bolin, B.B., Eriksson, E.E., 1959. Changes in the carbon dioxide content of the atmosphere and sea due to fossil fuel combustion. In: Bolin, B.B. (Ed.), *The Atmosphere and the Sea in Motion*. Scientific Contributions to the Rossby Memorial Volume. The Rockefeller Institute Press, New York, NY, pp. 130–142.
- Boudreau, B.P.B.P., 1997. *Diagenetic Models and Their Implementation*. vol. 505 Springer Berlin.
- Boudreau, B.P.B.P., Middelburg, J.J.J.J., Hofmann, A.F.A.F., Meysman, F.J.R.F.J.R., 2010. Ongoing transients in carbonate compensation. *Global Biogeochem. Cy.* 24. <http://dx.doi.org/10.1029/2009GB003654>. GB4010.
- Boudreau, B.P.B.P., Ruddick, B.R.B.R., 1991. On a reactive continuum representation of organic matter diagenesis. *Am. J. Sci.* 291 (5), 507–538.
- Bouttes, N.N., Paillard, D.D., Roche, D.D., 2010. Impact of brine-induced stratification on the glacial carbon cycle. *Clim. Past* 6, 575–589.
- Bouttes, N.N., Paillard, D.D., Roche, D.D., Waelbroeck, C.C., Kageyama, M.M., Laurantou, A.A., Michel, E.E., Bopp, L.L., 2012a. Impact of oceanic processes on the carbon cycle during the last termination. *Clim. Past* 8, 149–170.
- Bouttes, N.N., Paillard, D.D., Roche, D.D.M.D., Brovkin, V.V., Bopp, L.L., 2011. Last Glacial Maximum CO<sub>2</sub> and δ<sup>13</sup>C successfully reconciled. *Geophys. Res. Lett.* 38. <http://dx.doi.org/10.1029/2010GL044499>. <http://onlinelibrary.wiley.com/doi/10.1029/2010GL044499/abstract>.
- Bouttes, N.N., Roche, D.D.M.D., Paillard, D.D., 2009. Impact of strong deep ocean stratification on the glacial carbon cycle. *Paleoceanography* 24. <http://dx.doi.org/10.1029/2008PA001707>. <http://onlinelibrary.wiley.com/doi/10.1029/2008PA001707/abstract> PA3203.
- Bouttes, N.N., Roche, D.D.M.D., Paillard, D.D., 2012b. Systematic study of the impact of fresh water fluxes on the glacial carbon cycle. *Clim. Past* 8, 589–607. <http://dx.doi.org/10.5194/cp-8-589-2012>. <http://www.clim-past.net/8/589/2012/> 00007.
- Boyd, P.W.P.W., Trull, T.W.T.W., 2007. Understanding the export of biogenic particles in oceanic waters: is there consensus? *Prog. Oceanogr.* 72, 276–312. <http://dx.doi.org/10.1016/j.pocean.2006.10.007>. <http://www.sciencedirect.com/science/article/pii/S0079661106001340>.
- Bradley, J.A.J.A., Anesio, A.M.A.M., Arndt, S.S., 2016. Bridging the divide: a model-data approach to Polar and Alpine microbiology. *FEMS Microbiol. Ecol.* 92. <http://dx.doi.org/10.1093/femsec/fiw015>.
- Bralower, T.J.T.J., Meissner, K.J.K.J., Alexander, K.K., Thomas, D.J.D.J., 2014. The dynamics of global change at the Paleocene-Eocene thermal maximum: a data-model comparison. *Geochem. Geophys. Geosyst.* 15, 3830–3848. <http://dx.doi.org/10.1002/2014GC005474>. <http://onlinelibrary.wiley.com/doi/10.1002/2014GC005474/abstract> 00005.
- Broecker, W.W., 2003. The oceanic CaCO<sub>3</sub> cycle. *Treat. Geochem.* 6, 529–549.
- Broecker, W.S.W.S., Ledwell, J.R.J.R., Takahashi, T.T., Weiss, R.R., Merlivat, L.L., Memery, L.L., Peng, T.-H.T.-H., Jähne, B.B., Münnich, K.O.K.O., 1986. Isotopic versus micrometeorological ocean CO<sub>2</sub> fluxes: a serious conflict. *J. Geophys. Res.* 91, 10517–10527.
- Broecker, W.S.W.S., Peng, T.-H.T.-H., 1986. Carbon cycle: 1985. Glacial to interglacial changes in the operation of the global carbon cycle. *Radiocarbon* 28, 309–327.
- Broecker, W.S.W.S., Peng, T.-H.T.-H., 1989. The cause of the glacial to interglacial atmospheric CO<sub>2</sub> change: a polar alkalinity hypothesis. *Global Biogeochem. Cycles* 3, 215–239.
- Broecker, W.S.W.S., Peng, T.-H.T.-H., 1987. The role of CaCO<sub>3</sub> compensation in the glacial to interglacial atmospheric CO<sub>2</sub> change. *Global Biogeochem. Cycles* 1, 15–29.
- Broecker, W.S.W.S., Peng, T.-H.T.-H., Engh, R.R., 1980. Modeling the carbon system. *Radiocarbon* 22, 565–598. <http://dx.doi.org/10.1017/S003822200009966>.
- Broecker, W.S.W.S., Peng, T.-H.T.-H., Trumbore, S.S., Bonani, G.G., Wolfli, W.W., 1990. The distribution of radiocarbon in the glacial ocean. *Global Biogeochem. Cycles* 4, 103–117.
- Brovkin, V.V., Bendtsen, J.J., Claussen, M.M., Ganopolski, A.A., Kubatzki, C.C., Petoukhov, V.V., Andreev, A.A., 2002a. Carbon cycle, vegetation, and climate dynamics in the holocene: experiments with the CLIMBER-2 model. *Global Biogeochem. Cycles* 16, 1139. <http://dx.doi.org/10.1029/2001GB001662>. <http://onlinelibrary.wiley.com/doi/10.1029/2001GB001662/abstract>.
- Brovkin, V.V., Brücher, T.T., Kleinen, T.T., Zaehe, S.S., Joos, F.F., Roth, R.R., Spahni, R.R., Schmitt, J.J., Fischer, H.H., Leuenberger, M.M., Stone, E.J.E.J., Ridgwell, A.A., Chappellaz, J.J., Kehrwald, N.N., Barbante, C.C., Blunier, T.T., Dahl Jensen, D.D., 2016. Comparative carbon cycle dynamics of the present and last interglacial. *Quat. Sci. Rev.* 137, 15–32. <http://dx.doi.org/10.1016/j.quascirev.2016.01.028>. [http://www.sciencedirect.com/science/article/pii/S0273791163003000000000](http://www.sciencedirect.com/science/article/pii/S02737911630030000000).
- Brovkin, V.V., Ganopolski, A.A., Archer, D.D., Munhoven, G.G., 2012. Glacial CO<sub>2</sub> cycle as a succession of key physical and biogeochemical processes. *Clim. Past* 8, 251–264.
- Brovkin, V.V., Ganopolski, A.A., Archer, D.D., Rahmstorf, S.S., 2007. Lowering of glacial atmospheric CO<sub>2</sub> in response to changes in oceanic circulation and marine biogeochemistry. *Paleoceanography* 22. <http://dx.doi.org/10.1029/2006PA001380>. <http://onlinelibrary.wiley.com/doi/10.1029/2006PA001380/abstract> PA4202.
- Brovkin, V.V., Hofmann, M.M., Bendtsen, J.J., Ganopolski, A.A., 2002b. Ocean biology could control atmospheric δ<sup>13</sup>C during glacial-interglacial cycle. *Geochem. Geophys. Geosyst.* 3, 1–15. <http://dx.doi.org/10.1029/2001GC000270>. <http://onlinelibrary.wiley.com/doi/10.1029/2001GC000270/abstract>.
- Buesseler, K.O.K.O., Boyd, P.W.P.W., 2009. Shedding light on processes that control particle export and flux attenuation in the twilight zone of the open ocean. *Limnol. Oceanogr.* 54, 1210.
- Burdige, D.J.D.J., 2007. Preservation of organic matter in marine sediments: controls, mechanisms, and an imbalance in sediment organic carbon budgets? *Chem. Rev.* 107, 467–485. <http://dx.doi.org/10.1021/cr050347q>.
- Canfield, D.E.D.E., 1994. Factors influencing organic carbon preservation in marine sediments. *Chem. Geol.* 114, 315–329. [http://dx.doi.org/10.1016/0009-2541\(94\)90061-2](http://dx.doi.org/10.1016/0009-2541(94)90061-2). <http://www.sciencedirect.com/science/article/pii/0009254194900612>.
- Cao, L.L., Eby, M.M., Ridgwell, A.A., Caldeira, K.K., Archer, D.D., Ishida, A.A., Joos, F.F., Matsumoto, K.K., Mikolajewicz, U.U., Mouchet, A.A., Orr, J.C.J.C., Plattner, G.-K.G.-K., Schlitzer, R.R., Tokos, K.K., Tottterdell, I.I., Tschumi, T.T., Yamanaka, Y.Y., Yool, A.A., 2009. The role of ocean transport in the uptake of anthropogenic CO<sub>2</sub>.

- Biogeosciences 6, 375–390. <http://dx.doi.org/10.5194/bg-6-375-2009>.
- Cappellen, P.V.P.V., Wang, Y.Y., 1996. Cycling of iron and manganese in surface sediments: a general theory for the coupled transport and reaction of carbon, oxygen, nitrogen, sulfur, iron, and manganese. *Am. J. Sci.* 296, 197–243. <http://dx.doi.org/10.2475/ajs.296.3.197>. <http://www.ajsonline.org/content/296/3/197>.
- Chalikov, D.V.D.V., Verbitsky, M.Y.M.Y., 1984. A new earth climate model. *Nature* 308, 609–612. <http://dx.doi.org/10.1038/308609a0>. <http://www.nature.com/nature/journal/v308/n5960/abs/308609a0.html>.
- Ciais, P.P., Sabine, C.C., Bala, G.G., Bopp, L.L., Brovkin, V.V., Canadell, J.J., Chhabra, A.A., DeFries, R.R., Galloway, J.J., Heimann, M.M., et al., 2014. Carbon and other biogeochemical cycles. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, pp. 465–570.
- Claussen, M.M., Mysak, L.L., Weaver, A.A., Crucifix, M.M., Fichet, T.T., Loutre, M.-F.M.-F., Weber, S.S., Alcamo, J.J., Alexeev, V.V., Berger, A.A., Calov, R.R., Ganopolski, A.A., Goosse, H.H., Lohmann, G.G., Lunkeit, F.F., Mokhov, I.I., Petoukhov, V.V., Stone, P.P., Wang, Z.Z., 2002. Earth system models of intermediate complexity: closing the gap in the spectrum of climate system models. *Climate Dynam.* 18, 579–586. <http://dx.doi.org/10.1007/s00382-001-0200-1>. <http://link.springer.com/article/10.1007/s00382-001-0200-1>.
- Colbourn, G.G., Ridgwell, A.A., Lenton, T.M.T.M., 2013. The rock geochemical model (RokGeM) v0.9. *Geosci. Model Dev.* 6, 1543–1573. <http://dx.doi.org/10.5194/gmd-6-1543-2013>. <http://www.geosci-model-dev.net/6/1543/2013/>.
- Cox, P.M.P.M., Betts, R.A.R.A., Jones, C.D.C.D., Spall, S.A.S.A., Totterdell, I.J.I.J., 2000. Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model. *Nature* 408, 184–187. <http://dx.doi.org/10.1038/35041539>. <http://www.nature.com/nature/journal/v408/n6809/abs/408184a0.html>.
- Craig, H.H., 1957. The natural distribution of radiocarbon and the exchange time of carbon dioxide between atmosphere and sea. *Tellus* 9, 1–17. <http://dx.doi.org/10.1111/j.2153-3490.1957.tb01848.x>.
- Crucifix, M.M., 2005. Distribution of carbon isotopes in the glacial ocean: a model study. *Paleoceanography* 20. <http://dx.doi.org/10.1029/2005PA001131>. <http://onlinelibrary.wiley.com/doi/10.1029/2005PA001131/abstract> PA4020.
- Cui, Y.Y., Kump, L.L., Ridgwell, A.A., 2015. Spatial and Temporal Patterns of Ocean Acidification During the End-Permian Mass Extinction—An Earth System Model Evaluation, Volcanism and Global Environmental Change. Cambridge University Press, United Kingdom, pp. 291–306.
- Cui, Y.Y., Kump, L.R.L.R., Ridgwell, A.A., 2013. Initial assessment of the carbon emission rate and climatic consequences during the end-Permian mass extinction. *Paleoecol. Palaeoclimatol. Palaeoecol.* 389, 128–136. <http://dx.doi.org/10.1016/j.palaeo.2013.09.001>. <http://www.sciencedirect.com/science/article/pii/S0031018213003969>.
- Cui, Y.Y., Kump, L.R.L.R., Ridgwell, A.A., Charles, A.J.A.J., Junium, C.K.C.K., Diefendorf, A.F.A.F., Freeman, K.H.K.H., Urban, N.M.N.M., Harding, I.C.I.C., 2011. Slow release of fossil carbon during the Palaeocene-Eocene thermal maximum. *Nat. Geosci.* 4, 481–485. <http://dx.doi.org/10.1038/ngeo1179>. <http://www.nature.com/ngeo/journal/v4/n7/abs/ngeo1179.html>.
- Dalan, F.F., Stone, P.H.P.H., Kamenkovich, I.V.I.V., Scott, J.R.J.R., 2005. Sensitivity of the ocean's climate to diapycnal diffusivity in an EMC. Part I: equilibrium state. *J. Climate* 18, 2460–2481. <http://dx.doi.org/10.1175/JCLI3411.1>. <http://journals.ametsoc.org/doi/abs/10.1175/JCLI3411.1>.
- Dauwe, B.B., Middelburg, J.J.J.J., Herman, P.M.J.P.M.J., Heip, C.H.R.C.H.R., 1999. Linking diagenetic alteration of amino acids and bulk organic matter reactivity. *Limnol. Oceanogr.* 44, 1809–1814. <http://dx.doi.org/10.4319/lo.1999.44.7.1809>. [http://aslo.org/lo/toc/vol\\_44/issue\\_7/1809.html](http://aslo.org/lo/toc/vol_44/issue_7/1809.html).
- Davies, A.M.A.M., 1994. On the complementary nature of observational data, scientific understanding and model complexity: the need for a range of models. *J. Mar. Syst.* 5, 406–408. [http://dx.doi.org/10.1016/0924-7963\(94\)90003-5](http://dx.doi.org/10.1016/0924-7963(94)90003-5). <http://www.sciencedirect.com/science/article/pii/0924796394900035>.
- De La Rocha, C.L.C.L., Nowald, N.N., Passow, U.U., 2008. Interactions between diatom aggregates, minerals, particulate organic carbon, and dissolved organic matter: further implications for the ballast hypothesis. *Global Biogeochem. Cycles* 22. <http://dx.doi.org/10.1029/2007GB003156>. <http://onlinelibrary.wiley.com/doi/10.1029/2007GB003156/abstract>.
- Demaison, G.G., Moore, G.G., 1980. Anoxic environments and oil source bed genesis. *Org. Geochem.* 2, 9–31. [http://dx.doi.org/10.1016/0146-6380\(80\)90017-0](http://dx.doi.org/10.1016/0146-6380(80)90017-0). <http://www.sciencedirect.com/science/article/pii/0146638080900170>.
- Donnadieu, Y.Y., Godderis, Y.Y., Pierrehumbert, R.R., Dromart, G.G., Fluteau, F.F., Jacob, R.R., 2006. A GEOCLIM simulation of climatic and biogeochemical consequences of Pangea breakup. *Geochim. Geophys. Res.* 7. <http://dx.doi.org/10.1029/2006GC001278>. Q11019.
- Donnadieu, Y.Y., Ramstein, G.G., Goddés, Y.Y., Fluteau, F.F., 2004. Global tectonic setting and climate of the Late Neoproterozoic: a climate-geochemical coupled study. In: Jenkins, G.S.G.S., McMenamin, P.A.S.A., McKay, C.P.C.P., Sohl, L.L. (Eds.), *The Extreme Proterozoic: Geology, Geochemistry, and Climate*. American Geophysical Union, pp. 79–89. <http://onlinelibrary.wiley.com/doi/10.1029/146GM08/summary>.
- Ducklow, H.W.H.W., Quinby, H.L.L.L., Carlson, C.A.C.A., 1995. Bacterioplankton dynamics in the equatorial Pacific during the 1992 El Niño. *Deep-Sea Res.* II Top. Stud. Oceanogr. 42, 621–638. [http://dx.doi.org/10.1016/0967-0645\(95\)00022-1](http://dx.doi.org/10.1016/0967-0645(95)00022-1). <http://www.sciencedirect.com/science/article/pii/0967064595000221>.
- Dunne, J.P.J.P., Sarmiento, J.L.J.L., Gnanadesikan, A.A., 2007. A synthesis of global particle export from the surface ocean and cycling through the ocean interior and on the seafloor. *Global Biogeochem. Cycles* 21. <http://dx.doi.org/10.1029/2006GB002907>. <http://onlinelibrary.wiley.com/doi/10.1029/2006GB002907/abstract>.
- Duplessy, J.C.J.C., Roche, D.M.D.M., Kageyama, M.M., 2007. The deep ocean during the last interglacial period. *Science* 316, 89–91. <http://dx.doi.org/10.1126/science.1138582>. <http://www.sciencemag.org/content/316/5821/89>.
- Eby, M.M., Weaver, A.J.A.J., Alexander, K.K., Zickfeld, K.K., Abe-Ouchi, A.A., Cimattoribus, A.A.A.A., Crespin, E.E., Drijfhout, S.S.S.S., Edwards, N.R.N.R., Eliseev, A.V.A.V., Feulner, G.G., Fichet, T.T., Forest, C.E.C.E., Goosse, H.H., Holden, P.B.P.B., Joos, F.F., Kawamiya, M.M., Kicklighter, D.D., Kienert, H.H., Matsumoto, K.K., Mokhov, I.I.I.I., Monier, E.E., Olsen, S.M.S.M., Pedersen, J.O.P.J.O.P., Perrette, M.M., Philippon-Berthier, G.G., Ridgwell, A.A., Schlosser, A.A., Schneider von Deimling, T.T., Shaffer, G.G., Smith, R.S.R.S., Spahni, R.R., Sokolov, A.P.A.P., Steinacher, M.M., Tachiiri, K.K., Tokos, K.K., Yoshimori, M.M., Zeng, N.N., Zhao, F.F., 2013. Historical and idealized climate model experiments: an intercomparison of earth system models of intermediate complexity. *Clim. Past* 9, 1111–1140. <http://dx.doi.org/10.5194/cp-9-1111-2013>. <http://www.clim-past.net/9/1111/2013/>.
- Eby, M.M., Zickfeld, K.K., Montenegro, A.A., Archer, D.D., Meissner, K.J.K.J., Weaver, A.J.A.J., 2009. Lifetime of anthropogenic climate change: millennial time scales of potential CO<sub>2</sub> and surface temperature perturbations. *J. Climate* 22, 2501–2511. <http://dx.doi.org/10.1175/2008JCLI2554.1>. <http://journals.ametsoc.org/doi/abs/10.1175/2008JCLI2554.1>.
- Edwards, N.R.N.R., Marsh, R.R., 2005. Uncertainties due to transport-parameter sensitivity in an efficient 3-D ocean-climate model. *Climate Dynam.* 24, 415–433. <http://dx.doi.org/10.1007/s00382-004-0508-8>. <http://link.springer.com/article/10.1007/s00382-004-0508-8>.
- Emerson, S.S., Hedges, J.I.J.I., 1988. Processes controlling the organic carbon content of open ocean sediments. *Paleoceanography* 3, 621–634. <http://dx.doi.org/10.1029/PA003i005p0621>. <http://onlinelibrary.wiley.com/doi/10.1029/PA003i005p0621/abstract>.
- Eppley, R.W.R.W., Peterson, B.J.B.J., 1979. Particulate organic matter flux and planktonic new production in the deep ocean. *Nature* 282, 677–680.
- Eriksson, E.E., Welandar, P.P., 1956. On a mathematical model of the carbon cycle in nature. *Tellus* 8, 155–175. <http://dx.doi.org/10.1111/j.2153-3490.1956.tb01207.x>.
- Ewen, T.L.T.L., Weaver, A.J.A.J., Schmittner, A.A., 2004. Modelling carbon cycle feedbacks during abrupt climate change. *Quat. Sci. Rev.* 23, 431–448. <http://dx.doi.org/10.1016/j.quascirev.2003.08.007>. <http://www.sciencedirect.com/science/article/pii/S0273739103002889>.
- Fabry, V.J.V.J., 1989. Aragonite production by pteropod molluscs in the subarctic Pacific. *Deep Sea Res. Part A Oceanogr. Res.* 36, 1735–1751.
- Falkowski, P.P., Scholes, R.J.R.J., Boyle, E.E., Canadell, J.J., Canfield, D.D., Elser, J.J., Gruber, N.N., Hibbard, K.K., Höglberg, P.P., Linder, S.S., Mackenzie, F.T.F.T., Iii, B.M.B.M., Pedersen, T.T., Rosenthal, Y.Y., Seitzinger, S.S., Smetacek, V.V., Steffen, W.W., 2000. The global carbon cycle: a test of our knowledge of earth as a system. *Science* 290, 291–296. <http://dx.doi.org/10.1126/science.290.5490.291>. <http://www.sciencemag.org/content/290/5490/291>.
- Falkowski, P.G.P.G., Barber, R.T.R.T., Smetacek, V.V., 1998. Biogeochemical controls and feedbacks on ocean primary production. *Science* 281, 200–206. <http://dx.doi.org/10.1126/science.281.5374.200>. <http://www.sciencemag.org/content/281/5374/200>.
- Feely, R.A.R.A., Sabine, C.L.C.L., Lee, K.K., Berelson, W.W., Kleypas, J.J., Fabry, V.J.V.J., Millero, F.F.F.F., 2004. Impact of anthropogenic CO<sub>2</sub> on the CaCO<sub>3</sub> system in the oceans. *Science* 305, 362–366. <http://dx.doi.org/10.1126/science.1097329>. <http://www.sciencemag.org/content/305/5682/362>.
- Fiadeiro, M.M., 1980. The alkalinity of the deep Pacific. *Earth Planet. Sci. Lett.* 49, 499–505. [http://dx.doi.org/10.1016/0012-821X\(80\)90090-4](http://dx.doi.org/10.1016/0012-821X(80)90090-4). <http://www.sciencedirect.com/science/article/pii/0012821X80900904>.
- Frailé, I.I., Schulz, M.M., Mulitza, S.S., Merkel, U.U., Prange, M.M., Paul, A.A., 2009. Modeling the seasonal distribution of planktonic foraminifera during the Last Glacial Maximum. *Paleoceanography* 24. <http://dx.doi.org/10.1029/2008PA001686>. <http://onlinelibrary.wiley.com/doi/10.1029/2008PA001686/abstract> PA2216.
- Francois, R.R., Honjo, S.S., Krishfield, R.R., Manganini, S.S., 2002. Factors controlling the flux of organic carbon to the bathypelagic zone of the ocean. *Global Biogeochem. Cycles* 16. <http://dx.doi.org/10.1029/2001GB001722>. <http://onlinelibrary.wiley.com/doi/10.1029/2001GB001722/abstract> 34–1–34–20.
- Friedlingstein, P.P., Cox, P.P., Betts, R.R., Bopp, L.L., von Bloh, W.W., Brovkin, V.V., Cadule, P.P., Doney, S.S., Eby, M.M., Fung, I.I., Bala, G.G., John, J.J., Jones, C.C., Joos, F.F., Kato, T.T., Kawamiya, M.M., Knorr, W.W., Lindsay, K.K., Matthews, H.D.H.D., Raddatz, T.T., 2006. Climate-carbon cycle feedback analysis: results from the c4MIP model intercomparison. *J. Climate* 19, 3337–3353.
- Friedlingstein, P.P., Dufresne, J.-L.J.-L., Cox, P.M.P.M., Rayner, P.P., 2003. How positive is the feedback between climate change and the carbon cycle? *Tellus* B 55, 692–700. <http://dx.doi.org/10.1034/j.1600-0889.2003.01461.x>. <http://onlinelibrary.wiley.com/doi/10.1034/j.1600-0889.2003.01461.x/abstract>.
- Frieling, J.J., Svensen, H.H.H.H., Planke, S.S., Cramwinckel, M.J.M.J., Selnes, H.H., Sluijs, A.A., 2016. Thermogenic methane release as a cause for the long duration of the PETM. *Proc. Natl. Acad. Sci. USA* 113, 12059–12064. <http://dx.doi.org/10.1073/pnas.1603348113>.
- Galbraith, E.D.E.D., Kwon, E.Y.E.Y., Bianchi, D.D., Hain, M.P.M.P., Sarmiento, J.L.J.L., 2015. The impact of atmospheric pCO<sub>2</sub> on carbon isotope ratios of the atmosphere and ocean. *Global Biogeochem. Cycles* 29, 307–324. <http://dx.doi.org/10.1002/2014GB004929>.
- Angst, R.R., Joos, F.F., Gehlen, M.M., 2011. Sensitivity of pelagic calcification to ocean acidification. *Biogeosciences* 8, 433–458. <http://dx.doi.org/10.5194/bg-8-433-2011>. <http://www.biogeosciences.net/8/433/2011/>.
- Garrels, R.M.R.M., MacKenzie, F.T.F.T., 1972. A quantitative model for the sedimentary rock cycle. *Mar. Chem.* 1, 27–41. [http://dx.doi.org/10.1016/0304-4203\(72\)90004-7](http://dx.doi.org/10.1016/0304-4203(72)90004-7).
- Gehlen, M.M., Angst, R.R., Schneider, B.B., Bopp, L.L., Aumont, O.O., Ethe, C.C., 2007. The fate of pelagic CaCO<sub>3</sub> production in a high CO<sub>2</sub> ocean: a model study.

- Biogeosciences 4, 505–519.
- Gillooly, J.F.J.F., Brown, J.H.J.H., West, G.B.G.B., Savage, V.M.V.M., Charnov, E.L.E.L., 2001. Effects of size and temperature on metabolic rate. *Science* 293, 2248–2251. <http://dx.doi.org/10.1126/science.1061967>. <http://www.sciencemag.org/content/293/5538/2248>.
- Goddéris, Y.Y., Donnadié, Y.Y., de Vargas, C.C., Pierrehumbert, R.T.R.T., Dromart, G.G., van de Schootbrugge, B.B., 2008. Causal or casual link between the rise of nannoplankton calcification and a tectonically-driven massive decrease in late triassic atmospheric CO<sub>2</sub>? *Earth Planet. Sci. Lett.* 267, 247–255. <http://dx.doi.org/10.1016/j.epsl.2007.11.051>. <http://www.sciencedirect.com/science/article/pii/S0012821X07007753>.
- Goddéris, Y.Y., Joachimski, M.M., 2004. Global change in the Late Devonian: modelling the Frasnian-Famennian short-term carbon isotope excursions. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 202, 309–329. [http://dx.doi.org/10.1016/S0031-0182\(03\)00641-2](http://dx.doi.org/10.1016/S0031-0182(03)00641-2). <http://www.sciencedirect.com/science/article/pii/S0031018203006412>.
- Goodwin, P.P., Williams, R.G.R.G., Ridgwell, A.A., Follows, M.J.M.J., 2009. Climate sensitivity to the carbon cycle modulated by past and future changes in ocean chemistry. *Nat. Geosci.* 2, 145–150. <http://dx.doi.org/10.1038/ngeo416>. <http://www.nature.com/ngeo/journal/v2/n2/abs/ngeo416.html>.
- Goosse, H.H., Brovkin, V.V., Fichefet, T.T., Haarsma, R.R., Huybrechts, P.P., Jongma, J.J., Mouchet, A.A., Selten, F.F., Barriat, P.-Y.P.-Y., Campin, J.-M.J.-M., Deleersnijder, E.E., Driesschaert, E.E., Goelzer, H.H., Janssens, I.I., Loutre, M.-F.M.-F., Morales Maqueda, M.A.M.A., Opsteegh, T.T., Mathieu, P.-P.P.-P., Munhoven, G.G., Pettersson, E.J.E.J., Renssen, H.H., Roche, D.M.D.M., Schaeffer, M.M., Tartinville, B.B., Timmermann, A.A., Weber, S.L.S.L., 2010. Description of the earth system model of intermediate complexity LOVECLIM version 1.2. *Geosci. Model Dev.* 3, 603–633. <http://dx.doi.org/10.5194/gmd-3-603-2010>. <http://www.geosci-model-dev.net/3/603/2010/>.
- Gupta, N.S.N.S., Briggs, D.E.G.D.E.G., Collinson, M.E.M.E., Evershed, R.P.R.P., Michels, R.R., Jack, K.S.K.S., Pancost, R.D.R.D., 2007. Evidence for the in situ polymerisation of labile aliphatic organic compounds during the preservation of fossil leaves: implications for organic matter preservation. *Org. Geochem.* 38, 499–522. <http://dx.doi.org/10.1016/j.orgchem.2006.06.011>. <http://www.sciencedirect.com/science/article/pii/S0146638006001379>.
- Hain, M.P.M.P., Sigman, D.M.D.M., Haug, G.H.G.H., 2010. Carbon dioxide effects of Antarctic stratification, North Atlantic Intermediate Water formation, and sub-antarctic nutrient drawdown during the last ice age: diagnosis and synthesis in a geochemical box model. *Global Biogeochem. Cycles* 24, GB4023. <http://dx.doi.org/10.1029/2010GB003790>. <http://onlinelibrary.wiley.com/doi/10.1029/2010GB003790/abstract>.
- Hain, M.P.M.P., Sigman, D.M.D.M., Haug, G.H.G.H., 2014a. Distinct roles of the Southern Ocean and North Atlantic in the deglacial atmospheric radiocarbon decline. *Earth Planet. Sci. Lett.* 394, 198–208. <http://dx.doi.org/10.1016/j.epsl.2014.03.020>.
- Hain, M.P.M.P., Sigman, D.M.D.M., Haug, G.H.G.H., 2014b. The biological pump in the past. In: *Treatise on Geochemistry*, Second edition. Elsevier, Oxford, pp. 485–517. <http://www.sciencedirect.com/science/article/pii/B9780080959757006185>.
- Hales, B.B., 2003. Respiration, dissolution, and the lysocline. *Paleoceanography* 18, 1099. <http://dx.doi.org/10.1029/2003PA000915>. <http://onlinelibrary.wiley.com/doi/10.1029/2003PA000915/abstract>.
- Hales, B.B., Emerson, S.S., 1997. Evidence in support of first-order dissolution kinetics of calcite in seawater. *Earth Planet. Sci. Lett.* 148, 317–327. [http://dx.doi.org/10.1016/S0012-821X\(97\)00017-4](http://dx.doi.org/10.1016/S0012-821X(97)00017-4).
- Hansell, D.D., Carlson, C.C., Repeta, D.D., Schlitzer, R.R., 2009. Dissolved organic matter in the ocean: a controversy stimulates new insights. *Oceanography* 22, 202–211. <http://dx.doi.org/10.5670/oceanog.2009.109>. <https://dar.archive.mblwhoi.org/handle/1912/3183>.
- Hansell, D.A.D.A., 2013. Recalcitrant dissolved organic carbon fractions. *Ann. Rev. Mar. Sci.* 5, 421–445. <http://dx.doi.org/10.1146/annurev-marine-120710-100757>. <http://www.annualreviews.org/ep rint/hKnH2t8hM792BsymjTxB/full/10.1146/annurev-marine-120710-100757>.
- Hansell, D.A.D.A., Carlson, C.A.C.A., 2014. *Biogeochemistry of Marine Dissolved Organic Matter*. Academic Press.
- Hansell, D.A.D.A., Carlson, C.A.C.A., Schlitzer, R.R., 2012. Net removal of major marine dissolved organic carbon fractions in the subsurface ocean. *Global Biogeochem. Cycles* 26. <http://dx.doi.org/10.1029/2011GB004069>. <http://onlinelibrary.wiley.com/doi/10.1029/2011GB004069/abstract> GB1016.
- Hedges, J.I.J.I., Baldock, J.A.J.A., Gélinas, Y.Y., Lee, C.C., Peterson, M.M., Wakeham, S.G.S.G., 2001. Evidence for non-selective preservation of organic matter in sinking marine particles. *Nature* 409, 801–804. <http://dx.doi.org/10.1038/35057247>. <http://www.nature.com/nature/journal/v409/n6822/abs/409801a.html>.
- Hedges, J.I.J.I., Keil, R.G.R.G., 1995. Sedimentary organic matter preservation: an assessment and speculative synthesis. *Mar. Chem.* 49, 81–115. [http://dx.doi.org/10.1016/0304-4203\(95\)00008-F](http://dx.doi.org/10.1016/0304-4203(95)00008-F). <http://www.sciencedirect.com/science/article/pii/S030442039500008F>.
- Heine, C.C., Maier-Reimer, E.E., Winguth, A.M.A.M., Archer, D.D., 1999. A global oceanic sediment model for long-term climate studies. *Global Biogeochem. Cycles* 13, 221–250.
- Heine, M.M., Ilyina, T.T., 2015. Ocean biogeochemistry in the warm climate of the Late Paleocene. *Clim. Past* 11, 63–79. <http://dx.doi.org/10.5194/cp-11-63-2015>.
- Henson, S.A.S.A., Sanders, R.R., Madsen, E.E., 2012. Global patterns in efficiency of particulate organic carbon export and transfer to the deep ocean. *Global Biogeochem. Cycles* 26. <http://dx.doi.org/10.1029/2011GB004099>. <http://onlinelibrary.wiley.com/doi/10.1029/2011GB004099/abstract>.
- Henson, S.A.S.A., Sanders, R.R., Madsen, E.E., Morris, P.J.P.J., Le Moigne, F.F., Quartly, G.D.G.D., 2011. A reduced estimate of the strength of the ocean's biological carbon pump. *Geophys. Res. Lett.* 38. <http://dx.doi.org/10.1029/2011GL046735>. <http://onlinelibrary.wiley.com/doi/10.1029/2011GL046735/abstract> L04606 00050.
- Hoffert, M.I.M.I., Callegari, A.J.A.J., Hsieh, C.-T.C.-T., 1981. A box-diffusion carbon cycle model with upwelling, polar bottom water formation and a marine biosphere. In: Bolin, B.B. (Ed.), *Carbon Cycle Modelling*. SCOPE. vol. 16. John Wiley & Sons, Chichester, NY, pp. 285–305.
- Holden, P.P., Edwards, N.N., Müller, S.S., Oliver, K.K., Death, R.R., Ridgwell, A.A., 2013. Controls on the spatial distribution of oceanic  $\delta^{13}\text{C}$ . *Biogeosciences* 10, 1815–1833.
- Honjo, S.S., Manganini, S.J.S.J., Krishfield, R.A.R.A., Francois, R.R., 2008. Particulate organic carbon fluxes to the ocean interior and factors controlling the biological pump: a synthesis of global sediment trap programs since 1983. *Prog. Oceanogr.* 76, 217–285. <http://dx.doi.org/10.1016/j.pocan.2007.11.003>. <http://www.sciencedirect.com/science/article/pii/S0079661108000025>.
- Ibach, L.E.L.E.J., 1982. Relationship between sedimentation rate and total organic carbon content in ancient marine sediments. *AAPG Bulletin* 66, 170–188. <http://archives.datapages.com/data/bulletins/1982-83/data/pg/0066/0002/0150/0170.htm>.
- Iglesias-Rodriguez, M.D.M.D., Armstrong, R.R., Feely, R.R., Hood, R.R., Kleypas, J.J., Milliman, J.D.J.D., Sabine, C.C., Sarmiento, J.J., 2002. Progress made in study of ocean's calcium carbonate budget. *Eos, Trans. Am. Geophys. Union* 83, 365–375. <http://dx.doi.org/10.1029/2002EO000267>. <http://onlinelibrary.wiley.com/doi/10.1029/2002EO000267/abstract>.
- Jannasch, H.W.H.W., 1994. The microbial turnover of carbon in the deep-sea environment. *Global Planet. Change* 9, 289–295. [http://dx.doi.org/10.1016/0921-8181\(94\)90022-1](http://dx.doi.org/10.1016/0921-8181(94)90022-1). <http://www.sciencedirect.com/science/article/pii/0921818194900221>.
- Jansen, H.H., Zeebe, R.E.R.E., Wolf-Gladrow, D.A.D.A., 2002. Modeling the dissolution of settling CaCO<sub>3</sub> in the ocean. *Global Biogeochem. Cycles* 16. <http://dx.doi.org/10.1029/2000GB001279>. <http://onlinelibrary.wiley.com/doi/10.1029/2000GB001279/abstract> 11–1.
- Jiao, N.N., Herndl, G.J.G.J., Hansell, D.A.D.A., Benner, R.R., Kattner, G.G., Wilhelm, S.W.S.W., Kirchman, D.L.D.L., Weinbauer, M.G.M.G., Luo, T.T., Chen, F.F., Azam, F.F., 2010. Microbial production of recalcitrant dissolved organic matter: long-term carbon storage in the global ocean. *Nat. Rev. Microbiol.* 8, 593–599. <http://dx.doi.org/10.1038/nrmicro2386>. <http://www.nature.com/nrmicro/journal/v8/n8/abs/nrmicro2386.html>.
- John, E.H.E.H., Wilson, J.D.J.D., Pearson, P.N.P.N., Ridgwell, A.A., 2014. Temperature-dependent remineralization and carbon cycling in the warm Eocene oceans. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 413, 158–166. <http://dx.doi.org/10.1016/j.palaeo.2014.05.019>. <http://www.sciencedirect.com/science/article/pii/S0031018214002685>.
- Jones, N.S.N.S., Ridgwell, A.A., Hendy, E.J.E.J., 2015. Evaluation of coral reef carbonate production models at a global scale. *Biogeosciences* 12, 1339–1356. <http://dx.doi.org/10.5194/bg-12-1339-2015>. <http://www.biogeosciences.net/12/1339/2015/>.
- Jørgensen, B.B.B.B., 1978. A comparison of methods for the quantification of bacterial sulfate reduction in coastal marine sediments: II calculation from mathematical models. *Geomicrobiol. J.* 1, 29–47. <http://dx.doi.org/10.1080/01490457809377722>. <http://www.tandfonline.com/doi/abs/10.1080/01490457809377722>.
- Jourabchi, P.P., Cappellen, P.V.P.V., Regnier, P.P., 2005. Quantitative interpretation of pH distributions in aquatic sediments: a reaction-transport modeling approach. *Am. J. Sci.* 305, 919–956. <http://dx.doi.org/10.2475/ajs.305.9.919>. <http://www.ajsonline.org/content/305/9/919>.
- Kageyama, M.M., Braconnot, P.P., Harrison, S.P.S.P., Haywood, A.M.A.M., Jungclauss, J.J., Otto-Bliessner, B.L.B.L., Peterschmitt, J.-Y.J.-Y., Abe-Ouchi, A.A., Albani, S.S., Bartlein, P.J.P.J., Briery, C.C., Crucifix, M.M., Dolan, A.A., Fernandez-Donado, L.L., Fischer, H.H., Hopcroft, P.O.P.O., Ivanovic, R.F.R.F., Lambert, F.F., Lunt, D.J.D.J., Mahowald, N.M.N.M., Peltier, W.R.W.R., Phipps, S.J.S.J., Roche, D.M.D.M., Schmidt, G.A.G.A., Tarasov, L.L., Valdes, P.J.P.J., Zhang, Q.Q., Zhou, T.T., 2016. PMIP4-CMIP6: the contribution of the paleoclimate modelling intercomparison project to CMIP6. *Geosci. Model Dev. Discuss.* 2016, 1–46. <http://dx.doi.org/10.5194/gmd-2016-106>. <http://www.geosci-model-dev-discuss.net/gmd-2016-106/>.
- Keil, R.R., Hedges, J.J., 1993. Sorption of organic matter to mineral surfaces and the preservation of organic matter in coastal marine sediments. *Chem. Geol.* 107, 385–388.
- Keir, R.S.R.S., 1980. The dissolution kinetics of biogenic calcium carbonates in seawater. *Geochim. Cosmochim. Acta* 44, 241–252. [http://dx.doi.org/10.1016/0016-7037\(80\)90135-0](http://dx.doi.org/10.1016/0016-7037(80)90135-0). <http://www.sciencedirect.com/science/article/pii/0016703780901350>.
- Keir, R.S.R.S., 1988. On the Late Pleistocene ocean geochemistry and circulation. *Paleoceanography* 3, 413–445. <http://dx.doi.org/10.1029/PA003i004p00413>.
- Keir, R.S.R.S., 1990. Reconstructing the ocean carbon system variation during the last 150,000 years according to the Antarctic nutrient hypothesis. *Paleoceanography* 5, 253–276. <http://dx.doi.org/10.1029/PA005i003p00253>.
- Keir, R.S.R.S., 1991. The effect of vertical nutrient redistribution on surface ocean  $\delta^{13}\text{C}$ . *Global Biogeochem. Cycles* 5, 351–358. <http://dx.doi.org/10.1029/91GB01913>. <http://onlinelibrary.wiley.com/doi/10.1029/91GB01913/abstract>.
- Keir, R.S.R.S., Berger, W.H.W.H., 1983. Oceanic CO<sub>2</sub> content in the last 120,000 years: the phosphate-extraction model. *J. Geophys. Res.* 88, 6027–6038. <http://dx.doi.org/10.1029/JC088iC10p06027>.
- Keir, R.S.R.S., 1983. Reduction of thermohaline circulation during deglaciation: the effect on atmospheric radiocarbon and CO<sub>2</sub>. *Earth Planet. Sci. Lett.* 64, 445–456. [http://dx.doi.org/10.1016/0012-821X\(83\)90105-X](http://dx.doi.org/10.1016/0012-821X(83)90105-X).
- Keller, D.P.D.P., Oschlies, A.A., Eby, M.M., 2012. A new marine ecosystem model for the University of Victoria Earth System Climate Model. *Geosci. Model Dev.* 5, 1195–1220. <http://dx.doi.org/10.5194/gmd-5-1195-2012>. <http://www.geosci-model-dev.net/5/1195/2012/> 00018.
- Kirchman, D.L.D.L., Lancelot, C.C., Fasham, M.M., Legendre, L.L., Radach, G.G., Scott,

- M.M., 1993. Dissolved organic matter in biogeochemical models of the ocean. In: Evans, G.T.G.T., Fasham, M.J.R.M.J.R. (Eds.), *Towards a Model of Ocean Biogeochemical Processes*. no. 10 in NATO ASI Series. Springer Berlin Heidelberg. [http://link.springer.com/chapter/10.1007/978-3-642-84602-1\\_10](http://link.springer.com/chapter/10.1007/978-3-642-84602-1_10).
- Kirtland Turner, S.S., Ridgwell, A.A., 2013. Recovering the true size of an Eocene hyperthermal from the marine sedimentary record. *Paleoceanography* 28, 700–712. <http://dx.doi.org/10.1002/2013PA002541>. <http://onlinelibrary.wiley.com/doi/10.1002/2013PA002541/abstract>.
- Kirtland Turner, S.S., Ridgwell, A.A., 2016. Development of a novel empirical framework for interpreting geological carbon isotope excursions, with implications for the rate of carbon injection across the PETM. *Earth Planet. Sci. Lett.* 435, 1–13. <http://dx.doi.org/10.1016/j.epsl.2015.11.027>. <http://www.sciencedirect.com/science/article/pii/S0012821X15007311>.
- Klaas, C.C., Archer, D.E.D.E., 2002. Association of sinking organic matter with various types of mineral ballast in the deep sea: implications for the rain ratio. *Global Biogeochem. Cycles* 16 <http://dx.doi.org/10.1029/2001GB001765>. 63–1–63–14. <http://onlinelibrary.wiley.com/doi/10.1029/2001GB001765/abstract>.
- Kleinen, T.T., Brovkin, V.V., Munhoven, G.G., 2015. Carbon cycle dynamics during recent interglacials. *Climate Past Discuss.* 11, 1945–1983. <http://dx.doi.org/10.5194/cpd-11-1945-2015>. <http://www.clim-past-discuss.net/11/1945/2015/>.
- Kleydas, J.A.J.A., Langdon, C.C., 2006. Coral reefs and changing seawater carbonate chemistry. *Coast. Estuar. Stud. Coral Reefs Clim. Chang. Sci. Manag.* 61, 73–110.
- Knox, F.F., McElroy, M.B.M.B., 1984. Changes in atmospheric CO<sub>2</sub>: influence of the marine biota at high latitude. *J. Geophys. Res.* 89, 4629–4637. <http://dx.doi.org/10.1029/JD089iD03p04629>.
- Kohfeld, K.E.K.E., Quéré, C.L.C.L., Harrison, S.P.S.P., Anderson, R.F.R.F., 2005. Role of marine biology in glacial-interglacial CO<sub>2</sub> cycles. *Science* 308, 74–78. <http://dx.doi.org/10.1126/science.1105375>. <http://science.sciencemag.org/content/308/5718/74>.
- Köhler, P.P., 2016. Using the Suess effect on the stable carbon isotope to distinguish the future from the past in radiocarbon. *Environ. Res. Lett.* 11, 124016. <http://dx.doi.org/10.1088/1748-9326/11/12/124016>.
- Köhler, P.P., Bintanja, R.R., 2008. The carbon cycle during the mid pleistocene transition: the southern ocean decoupling hypothesis. *Clim. Past* 4, 311–332.
- Köhler, P.P., Fischer, H.H., 2006. Simulating low frequency changes in atmospheric CO<sub>2</sub> during the last 740 000 years. *Clim. Past* 2, 57–78.
- Köhler, P.P., Fischer, H.H., Munhoven, G.G., Zeebe, R.E.R.E., 2005. Quantitative interpretation of atmospheric carbon records over the last glacial termination. *Global Biogeochem. Cycles* 19 <http://dx.doi.org/10.1029/2004GB002345>. <http://onlinelibrary.wiley.com/doi/10.1029/2004GB002345/abstract>.
- Köhler, P.P., Fischer, H.H., Schmitt, J.J., 2010. Atmospheric δ<sup>13</sup>C<sub>CO<sub>2</sub> and its relation to pCO<sub>2</sub> and deep ocean δ<sup>13</sup>C during the Late Pleistocene. *Paleoceanography* 25, PA1213. <http://dx.doi.org/10.1029/2008PA001703>. <http://onlinelibrary.wiley.com/doi/10.1029/2008PA001703/abstract>.</sub>
- Köhler, P.P., Muscheler, R.R., Fischer, H.H., 2006. A model-based interpretation of low-frequency changes in the carbon cycle during the last 120,000 years and its implications for the reconstruction of atmospheric delta C-14. *Geochem. Geophys. Geosyst.* 7 <http://dx.doi.org/10.1029/2005GC001228>. Q11N06.
- Komar, N.N., Zeebe, R.E.R.E., 2011. Oceanic calcium changes from enhanced weathering during the Paleocene-Eocene thermal maximum: no effect on calcium-based proxies. *Paleoceanography* 26 <http://dx.doi.org/10.1029/2010PA001979>. <http://onlinelibrary.wiley.com/doi/10.1029/2010PA001979/abstract> PA3211.
- Komar, N.N., Zeebe, R.E.R.E., Dickens, G.R.G.R., 2013. Understanding long-term carbon cycle trends: the Late Paleocene through the Early Eocene. *Paleoceanography* 28, 650–662. <http://dx.doi.org/10.1002/palo.20060>. <http://onlinelibrary.wiley.com/doi/10.1002/palo.20060/abstract>.
- Kriest, I.I., Khatiwala, S.S., Oshlies, A.A., 2010. Towards an assessment of simple global marine biogeochemical models of different complexity. *Prog. Oceanogr.* 86, 337–360. <http://dx.doi.org/10.1016/j.pocan.2010.05.002>. <http://www.sciencedirect.com/science/article/pii/S0079661110000844>.
- Kriest, I.I., Oshlies, A.A., 2013. Swept under the carpet: organic matter burial decreases global ocean biogeochemical model sensitivity to remineralization length scale. *Biogeosciences* 10, 8401–8422. <http://dx.doi.org/10.5194/bg-10-8401-2013>. <http://www.biogeosciences.net/10/8401/2013/>.
- Kriest, I.I., Oshlies, A.A., Khatiwala, S.S., 2012. Sensitivity analysis of simple global marine biogeochemical models. *Global Biogeochem. Cycles* 26, GB2029. <http://dx.doi.org/10.1029/2011GB004072>. <http://onlinelibrary.wiley.com/doi/10.1029/2011GB004072/abstract>.
- Kujawinski, E.B.E.B., 2011. The impact of microbial metabolism on marine dissolved organic matter. *Ann. Rev. Mar. Sci.* 3, 567–599. <http://dx.doi.org/10.1146/annurev-marine-120308-081003>. <http://dx.doi.org/10.1146/annurev-marine-120308-081003>.
- Kump, L.R.L.R., Brantley, S.L.S.L., Arthur, M.A.M.A., 2000. Chemical weathering, atmospheric CO<sub>2</sub>, and climate. *Annu. Rev. Earth Planet. Sci.* 28, 611–667. <http://dx.doi.org/10.1146/annurev.earth.28.1.611>. <http://www.annualreviews.org/doi/abs/10.1146/annurev.earth.28.1.611>.
- Kwon, E.Y.E.Y., Primeau, F.F., Sarmiento, J.L.J.L., 2009. The impact of remineralization depth on the air-sea carbon balance. *Nat. Geosci.* 2, 630–635. <http://dx.doi.org/10.1038/ngeo612>. <http://www.nature.com/ngeo/journal/v2/n9/abs/ngeo612.html>.
- Lam, P.J.P.J., Doney, S.C.S.C., Bishop, J.K.B.J.K.B., 2011. The dynamic ocean biological pump: insights from a global compilation of particulate organic carbon, CaCO<sub>3</sub>, and opal concentration profiles from the mesopelagic. *Global Biogeochem. Cycles* 25 <http://dx.doi.org/10.1029/2010GB003868>. <http://onlinelibrary.wiley.com/doi/10.1029/2010GB003868/abstract>.
- Lambert, F.F., Tagliabue, A.A., Shaffer, G.G., Lamy, F.F., Winckler, G.G., Farias, L.L., Gallardo, L.L., De Pol-Holz, R.R., 2015. Dust fluxes and iron fertilization in Holocene and Last Glacial Maximum climates. *Geophys. Res. Lett.* 42 <http://dx.doi.org/10.1002/2015GL064250>. <http://onlinelibrary.wiley.com/doi/10.1002/2015GL064250/abstract> 2015GL064250.
- Lane, E.E., Peacock, S.S., Restrepo, J.M.J.M., 2006. A dynamic-flow carbon-cycle box model and high-latitude sensitivity. *Tellus* 58B, 257–278. <http://dx.doi.org/10.1111/j.1600-0889.2006.00192.x>.
- Lasaga, A.C.A.C., 1981. Rate laws of chemical reactions. In: Lasaga, A.A., Kirkpatrick, R.J.R.J. (Eds.), *Reviews in Mineralogy*: (United States). 8. pp. 1–68. <http://www.osti.gov/scitech/biblio/6034791>.
- Laws, E.A.E.A., Falkowski, P.G.P.G., Smith, W.O.W.O., Ducklow, H.H., McCarthy, J.J.J.J., 2000. Temperature effects on export production in the open ocean. *Global Biogeochem. Cycles* 14, 1231–1246. <http://dx.doi.org/10.1029/1999GB001229>. <http://onlinelibrary.wiley.com/doi/10.1029/1999GB001229/abstract>.
- Le Hir, G.G., Goddérès, Y.Y., Donnadiou, Y.Y., Ramstein, G.G., 2008. A geochemical modelling study of the evolution of the chemical composition of seawater linked to a “snowball” glaciation. *Biogeosciences* 5, 253–267. <http://hal.archives-ouvertes.fr/hal-00330345>.
- Le Moigne, F.A.C.F.A.C., Pabortsava, K.K., Marcinko, C.L.J.C.L.J., Martin, P.P., Sanders, R.J.R.J., 2014. Where is mineral ballast important for surface export of particulate organic carbon in the ocean? *Geophys. Res. Lett.* 41 <http://dx.doi.org/10.1002/2014GL061678>. <http://onlinelibrary.wiley.com/doi/10.1002/2014GL061678/abstract> 2014GL061678.
- Le Moigne, F.A.C.F.A.C., Sanders, R.J.R.J., Villa-Alfageme, M.M., Martin, A.P.A.P., Pabortsava, K.K., Planquette, H.H., Morris, P.J.P.J., Thomalla, S.J.S.J., 2012. On the proportion of ballast versus non-ballast associated carbon export in the surface ocean. *Geophys. Res. Lett.* 39, L15610. <http://dx.doi.org/10.1029/2012GL052980>. <http://onlinelibrary.wiley.com/doi/10.1029/2012GL052980/abstract>.
- Locarnini, R.R., Mishonov, A.A., Antonov, J.J., Boyer, T.T., Garcia, H.H., Baranova, O.O., Zweng, M.M., Paver, C.C., Reagan, J.J., Johnson, D.D., et al., 2013. *World Ocean Atlas 2013*. Vol. 1: Temperature. A. Mishonov, Technical Ed. NOAA Atlas NESDIS, 73, 40. pp. 40.
- Logan, B.E.B.E., Hunt, J.R.J.R., 1987. Advantages to microbes of growth in permeable aggregates in marine systems. *Limnol. Oceanogr.* 32, 1034–1048.
- López-Urrutia, A.A., Martin, E.S.E.S., Harris, R.P.R.P., Irigoien, X.X., 2006. Scaling the metabolic balance of the oceans. *Proc. Natl. Acad. Sci.* 103, 8739–8744. <http://dx.doi.org/10.1073/pnas.0601137103>. <http://www.pnas.org/content/103/23/8739>.
- Lunt, D.J.D.J., Huber, M.M., Anagnostou, E.E., Baatsen, M.L.M.L.J., Caballero, R.R., DeConto, R.R., Dijkstra, H.A.H.A., Donnadiou, Y.Y., Evans, D.D., Feng, R.R., Foster, G.L.G.L., Gasson, E.E., von der Heydt, A.S.A.S., Hollis, C.J.C.J., Inglis, G.N.G.N., Jones, S.S.M.M., Kiehl, J.J., Kirtland Turner, S.S., Korty, R.L.R.L., Kozyro, R.R., Krishnan, S.S., Ladant, J.-B.J.-B., Langebroek, P.P., Lear, C.H.C.H., LeGrande, A.N.A.N., Littler, K.K., Markwick, P.P., Otto-Bliesner, B.B., Pearson, P.P., Poulsen, C.J.C.J., Salzman, U.U., Shields, C.C., Snell, K.K., Stärr, M.M., Super, J.J., Tabor, C.C., Tierney, J.E.J.E., Tourre, G.J.L.G.J.L., Tripati, A.A., Upchurch, G.R.G.R., Wade, B.S.B.S., Wing, S.L.S.L., Winguth, A.M.E.A.M.E., Wright, N.M.N.M., Zachos, J.C.J.C., Zeebe, R.E.R.E., 2017. The deepMIP contribution to PMIP4: experimental design for model simulations of the EECO, PETM, and pre-PETM (version 1.0). *Geosci. Model Dev.* 10, 889–901. <http://dx.doi.org/10.5194/gmd-10-889-2017>. <http://www.geosci-model-dev.net/10/889/2017/>.
- Lutz, M.M., Dunbar, R.R., Caldeira, K.K., 2002. Regional variability in the vertical flux of particulate organic carbon in the ocean interior. *Global Biogeochem. Cycles* 16 <http://dx.doi.org/10.1029/2000GB001383>. <http://onlinelibrary.wiley.com/doi/10.1029/2000GB001383/abstract> 11–1–11–18.
- Ma, W.W., Tian, J.J., 2014. Modeling the contribution of dissolved organic carbon to carbon sequestration during the Last Glacial Maximum. *Geo-Mar. Lett.* 34, 471–482. <http://dx.doi.org/10.1007/s00367-014-0378-y>. <http://link.springer.com/article/10.1007/s00367-014-0378-y>.
- Ma, Z.Z., Gray, E.E., Thomas, E.E., Murphy, B.B., Zachos, J.J., Paytan, A.A., 2014. Carbon sequestration during the Palaeocene-Eocene Thermal Maximum by an efficient biological pump. *Nat. Geosci.* 7, 382–388. <http://dx.doi.org/10.1038/ngeo2139>. <http://www.nature.com/ngeo/journal/v7/n5/full/ngeo2139.html>.
- Mackenzie, F.T.F.T., Lerman, A.A., Andersson, A.J.A.J., 2004. Past and present of sediment and carbon biogeochemical cycling models. *Biogeosciences* 1, 11–32.
- Maier-Reimer, E.E., 1993. Geochemical cycles in an ocean general circulation model. Preindustrial tracer distributions. *Global Biogeochem. Cycles* 7, 645–677. <http://dx.doi.org/10.1029/93GB01355>. <http://onlinelibrary.wiley.com/doi/10.1029/93GB01355/abstract>.
- Maier-Reimer, E.E., Hasselmann, K.K., 1987. Transport and storage of CO<sub>2</sub> in the ocean – an inorganic ocean-circulation carbon cycle model. *Clim. Dynam.* 2, 63–90. <http://dx.doi.org/10.1007/BF01054491>.
- Marchal, O.O., Stocker, T.F.T.F., Joos, F.F., 1998a. Impact of oceanic reorganizations on the ocean carbon cycle and atmospheric carbon dioxide content. *Paleoceanography* 13, 225–244. <http://dx.doi.org/10.1029/98PA00726>. <http://onlinelibrary.wiley.com/doi/10.1029/98PA00726/abstract>.
- Marchal, O.O., Stocker, T.F.T.F., Joos, F.F., 1998b. A latitude-depth, circulation-biogeophysical ocean model for paleoclimatic studies. Development and sensitivities. *Tellus* B 50, 290–316. <http://dx.doi.org/10.1034/j.1600-0889.1998.t01-2-00006.x>. <http://onlinelibrary.wiley.com/doi/10.1034/j.1600-0889.1998.t01-2-00006.x/abstract>.
- Marchal, O.O., Stocker, T.F.T.F., Joos, F.F., 1999a. Physical and biogeochemical responses to freshwater-induced thermohaline variability in a zonally averaged ocean model. In: Clark, U.U., Webb, S.S., Keigwin, D.D. (Eds.), *Geophysical Monograph Series*. vol. 112. American Geophysical Union, Washington, D. C., pp. 263–284. <http://www.agu.org/books/gm/v112/GM112p0263/GM112p0263.shtml>.
- Marchal, O.O., Stocker, T.F.T.F., Joos, F.F., Indermühle, A.A., Blunier, T.T., Tschumi, J.J., 1999b. Modelling the concentration of atmospheric CO<sub>2</sub> during the younger dryas climate event. *Climate Dynam.* 15, 341–354. <http://dx.doi.org/10.1007/>

- s003820050286. <http://link.springer.com/article/10.1007/s003820050286>.
- Marchal, O.O., Stocker, T.F.T.F., Muscheler, R.R., 2001. Atmospheric radiocarbon during the younger dryas: production, ventilation, or both? *Earth Planet. Sci. Lett.* 185, 383–395. [http://dx.doi.org/10.1016/S0012-821X\(00\)00383-6](http://dx.doi.org/10.1016/S0012-821X(00)00383-6). <http://www.sciencedirect.com/science/article/pii/S0012821X00003836>.
- Marinov, I.I., Gnanadesikan, A.A., Sarmiento, J.L.J.L., Toggweiler, J.R.J.R., Follows, M.M., Mignone, B.K.B.K., 2008. Impact of oceanic circulation on biological carbon storage in the ocean and atmospheric pCO<sub>2</sub>. *Global Biogeochem. Cycles* 22, GB3007. <http://dx.doi.org/10.1029/2007GB002958>. <http://onlinelibrary.wiley.com/doi/10.1029/2007GB002958/abstract>.
- Marsay, C.M.C.M., Sanders, R.J.R.J., Henson, S.A.S.A., Pabortsava, K.K., Achterberg, E.P.E.P., Lampitt, R.S.R.S., 2015. Attenuation of sinking particulate organic carbon flux through the mesopelagic ocean. *Proc. Natl. Acad. Sci.* 112, 1089–1094. <http://dx.doi.org/10.1073/pnas.1415311112>. <http://www.pnas.org/content/112/4/1089>.
- Martin, J.H.J.H., Fitzwater, S.E.S.E., Michael Gordon, R.R., Hunter, C.N.C.N., Tanner, S.J.S.J., 1993. Iron, primary production and carbon-nitrogen flux studies during the JGOFS North Atlantic bloom experiment. *Deep-Sea Res. II Top. Stud. Oceanogr.* 40, 115–134. [http://dx.doi.org/10.1016/0967-0645\(93\)90009-C](http://dx.doi.org/10.1016/0967-0645(93)90009-C). <http://www.sciencedirect.com/science/article/pii/096706459390009C> 00276.
- Martin, J.H.J.H., Knauer, G.A.G.A., Karl, D.M.D.M., Broenkow, W.W.W.W., 1987. VERTEX: carbon cycling in the northeast Pacific. *Deep Sea Res. Part A Oceanogr. Res. Pap.* 34, 267–285. [http://dx.doi.org/10.1016/0198-0149\(87\)90086-0](http://dx.doi.org/10.1016/0198-0149(87)90086-0). <http://www.sciencedirect.com/science/article/pii/0198014987900860>.
- Matsumoto, K.K., 2007. Biology-mediated temperature control on atmospheric pCO<sub>2</sub> and ocean biogeochemistry. *Geophys. Res. Lett.* 34, L20605. <http://dx.doi.org/10.1029/2007GL031301>. <http://onlinelibrary.wiley.com/doi/10.1029/2007GL031301/abstract>.
- Matsumoto, K.K., Yokoyama, Y.Y., 2013. Atmospheric Δ<sup>14</sup>C reduction in simulations of Atlantic overturning circulation shutdown. *Glob. Biogeochem. Cycles* 27, 296–304. <http://dx.doi.org/10.1002/gbc.20035>. <http://onlinelibrary.wiley.com/doi/10.1002/gbc.20035/abstract>.
- Matsumoto, K.K., Chase, Z.Z., Kohfeld, K.K., 2014. Different mechanisms of silicic acid leakage and their biogeochemical consequences. *Paleoceanography* 29. <http://dx.doi.org/10.1002/2013PA002588>. <http://onlinelibrary.wiley.com/doi/10.1002/2013PA002588/abstract> 2013PA002588.
- Matsumoto, K.K., Tokos, K.S.K.S., Price, A.R.A.R., Cox, S.J.S.J., 2008. First description of the Minnesota Earth System Model for Ocean biogeochemistry (MESMO 1.0). *Geosci. Model Dev.* 1, 1–15. <http://dx.doi.org/10.5194/gmd-1-1-2008>. <http://www.geosci-model-dev.net/1/1/2008/>.
- McGuffie, K.K., Henderson-Sellers, A.A., 2005. *A Climate Modelling Primer*. John Wiley & Sons.
- Meinshausen, M.M., Raper, S.C.B.S.C.B., Wigley, T.M.L.T.M.L., 2011. Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 - part 1: model description and calibration. *Atmos. Chem. Phys.* 11, 1417–1456. <http://dx.doi.org/10.5194/acp-11-1417-2011>.
- Meissner, K.J.K.J., Bralower, T.J.T.J., Alexander, K.K., Jones, T.D.T.D., Sijp, W.W., Ward, M.M., 2014. The Paleocene-Eocene Thermal Maximum: how much carbon is enough? *Paleoceanography* 29, 946–963. <http://dx.doi.org/10.1002/2014PA002650>. <http://onlinelibrary.wiley.com/doi/10.1002/2014PA002650/abstract> 00003.
- Melesse, A.M.A.M., Hanley, R.S.R.S., 2005. Artificial neural network application for multi-ecosystem carbon flux simulation. *Ecol. Model.* 189, 305–314. <http://dx.doi.org/10.1016/j.ecolmodel.2005.03.014>. <http://www.sciencedirect.com/science/article/pii/S0304380005002061> 00070.
- Menviel, L.L., Joos, F.F., 2012. Toward explaining the holocene carbon dioxide and carbon isotope records: results from transient ocean carbon cycle-climate simulations. *Paleoceanography* 27. <http://dx.doi.org/10.1029/2011PA002224>. <http://dx.doi.org/10.1029/2011PA002224> PA1207.
- Menviel, L.L., England, M.H.M.H., Meissner, K.J.K.J., Mouchet, A.A., Yu, J.J., 2014. Atlantic-Pacific seesaw and its role in outgassing CO<sub>2</sub> during Heinrich events. *Paleoceanography* 29, 58–70. <http://dx.doi.org/10.1002/2013PA002542>. <http://onlinelibrary.wiley.com/doi/10.1002/2013PA002542/abstract> 00018.
- Menviel, L.L., Joos, F.F., Ritz, S.S., 2012. Simulating atmospheric CO<sub>2</sub>, <sup>13</sup>C and the marine carbon cycle during the Last Glacial-Interglacial cycle: possible role for a deepening of the mean remineralization depth and an increase in the oceanic nutrient inventory. *Quat. Sci. Rev.* 56, 46–68. <http://dx.doi.org/10.1016/j.quascirev.2012.09.012>. <http://www.sciencedirect.com/science/article/pii/S0277379112003496>.
- Menviel, L.L., Timmermann, A.A., Mouchet, A.A., Timm, O.O., 2008a. Climate and marine carbon cycle response to changes in the strength of the Southern Hemispheric westerlies. *Paleoceanography* 23. <http://dx.doi.org/10.1029/2008PA001604>. <http://onlinelibrary.wiley.com/doi/10.1029/2008PA001604/abstract> PA4201.
- Menviel, L.L., Timmermann, A.A., Mouchet, A.A., Timm, O.O., 2008b. Meridional reorganizations of marine and terrestrial productivity during Heinrich events. *Paleoceanography* 23. <http://dx.doi.org/10.1029/2007PA001445>. PA1203.
- Menviel, L.L., Timmermann, A.A., Timm, O.E.O.E., Mouchet, A.A., 2010. Climate and biogeochemical response to a rapid melting of the West Antarctic Ice Sheet during interglacials and implications for future climate. *Paleoceanography* 25. <http://dx.doi.org/10.1029/2009PA001892>. <http://onlinelibrary.wiley.com/doi/10.1029/2009PA001892/abstract> PA4231.
- Menviel, L.L., Timmermann, A.A., Timm, O.E.O.E., Mouchet, A.A., 2011. Deconstructing the last glacial termination: the role of millennial and orbital-scale forcings. *Quat. Sci. Rev.* 30, 1155–1172. <http://dx.doi.org/10.1016/j.quascirev.2011.02.005>. <http://www.sciencedirect.com/science/article/pii/S0277379111000539>.
- Meyer, K.M.K.M., Kump, L.R.L.R., 2008. Oceanic euxinia in Earth history: causes and consequences. *Annu. Rev. Earth Planet. Sci.* 36, 251–288. <http://dx.doi.org/10.1146/annurev.earth.36.031207.124256>. <http://dx.doi.org/10.1146/annurev.earth.36.031207.124256> 00175.
- Meyer, K.M.K.M., Kump, L.R.L.R., Ridgwell, A.A., 2008. Biogeochemical controls on photic-zone euxinia during the end-Permian mass extinction. *Geology* 36, 747–750. <http://dx.doi.org/10.1130/G24618A.1>. <http://geology.gsapubs.org/content/36/9/747>.
- Meyers, S.R.S.R., 2007. Production and preservation of organic matter: the significance of iron. *Paleoceanography* 22. <http://dx.doi.org/10.1029/2006PA001332>. <http://onlinelibrary.wiley.com/doi/10.1029/2006PA001332/abstract> PA4211.
- Michel, E.E., Labeyrie, L.D.L.D., Duplessy, J.-C.J.-C., Gorfli, N.N., Labracherie, M.M., Turon, J.-L.J.-L., 1995. Could deep subantarctic convection feed the world deep basins during the Last Glacial Maximum? *Paleoceanography* 10, 927–941. <http://dx.doi.org/10.1029/95PA00978>.
- Middelburg, J.J.J.J., 1989. A simple rate model for organic matter decomposition in marine sediments. *Geochim. Cosmochim. Acta* 53, 1577–1581. [http://dx.doi.org/10.1016/0016-7037\(89\)90239-1](http://dx.doi.org/10.1016/0016-7037(89)90239-1). <http://www.sciencedirect.com/science/article/pii/0016703789902391>.
- Middelburg, J.J.J.J., Meysman, F.J.R.F.J.R., 2007. Burial at sea. *Science* 316, 1294–1295. <http://dx.doi.org/10.1126/science.1144001>. <http://www.vliz.be/IMIS/imis.php?module=ref&refid=111049>.
- Millero, F.F.F.F., 2007. The marine inorganic carbon cycle. *Chem. Rev.* 107, 308–341. <http://dx.doi.org/10.1021/cr0503557>.
- Milliman, J.D.J.D., 1993. Production and accumulation of calcium carbonate in the ocean: budget of a nonsteady state. *Global Biogeochem. Cycles* 7, 927–957. <http://dx.doi.org/10.1029/93GB02524>. <http://onlinelibrary.wiley.com/doi/10.1029/93GB02524/abstract>.
- Milliman, J.D.J.D., Droxler, A.W.A.W., 1996. Neritic and pelagic carbonate sedimentation in the marine environment: ignorance is not bliss. *Geol. Rundsch.* 85, 496–504. <http://dx.doi.org/10.1007/BF02369004>. <http://link.springer.com/article/10.1007/BF02369004>.
- Milliman, J.D.J.D., Troy, P.J.P.J., Balch, W.M.W.M., Adams, A.K.A.K., Li, Y.H.Y.H., Mackenzie, F.T.F.T., 1999. Biologically mediated dissolution of calcium carbonate above the chemical lysocline? *Deep-Sea Res. I Oceanogr. Res. Pap.* 46, 1653–1669. [http://dx.doi.org/10.1016/S0967-0637\(99\)00034-5](http://dx.doi.org/10.1016/S0967-0637(99)00034-5). <http://www.sciencedirect.com/science/article/pii/S0967063799000345>.
- Monteiro, F.M.F.M., Pancost, R.D.R.D., Ridgwell, A.A., Donnadieu, Y.Y., 2012. Nutrients as the dominant control on the spread of anoxia and euxinia across the Cenomanian-Turonian oceanic anoxic event (OAE2): model-data comparison. *Paleoceanography* 27. <http://dx.doi.org/10.1029/2012PA002351>. <http://onlinelibrary.wiley.com/doi/10.1029/2012PA002351/abstract>.
- Montenegro, A.A., Spence, P.P., Meissner, K.J.K.J., Eby, M.M., Melchin, M.J.M.J., Johnston, S.T.S.T., 2011. Climate simulations of the Permian-Triassic boundary: ocean acidification and the extinction event. *Paleoceanography* 26, PA3207. <http://dx.doi.org/10.1029/2010PA002058>. <http://onlinelibrary.wiley.com/doi/10.1029/2010PA002058/abstract> 00014.
- Morse, J.W.J.W., Arvidson, R.S.R.S., 2002. The dissolution kinetics of major sedimentary carbonate minerals. *Earth Sci. Rev.* 58, 51–84. [http://dx.doi.org/10.1016/S0012-8252\(01\)00083-6](http://dx.doi.org/10.1016/S0012-8252(01)00083-6). <http://www.sciencedirect.com/science/article/pii/S0012825201000836>.
- Morse, J.W.J.W., Arvidson, R.S.R.S., Lüttge, A.A., 2007. Calcium carbonate formation and dissolution. *Chem. Rev.* 107, 342–381. <http://dx.doi.org/10.1021/cr050358j>.
- Morse, J.W.J.W., Berner, R.A.R.A., 1972. Dissolution kinetics of calcium carbonate in sea water; I, a kinetic origin for the lysocline. *Am. J. Sci.* 272, 840–851. <http://dx.doi.org/10.2475/ajs.272.9.840>. <http://www.ajsonline.org/content/272/9/840>.
- Morse, J.W.J.W., He, S.S., 1993. Influences of T, S and pCO<sub>2</sub> on the pseudo-homogeneous precipitation of CaCO<sub>3</sub> from seawater: implications for whitening formation. *Mar. Chem.* 41, 291–297. [http://dx.doi.org/10.1016/0304-4203\(93\)90261-L](http://dx.doi.org/10.1016/0304-4203(93)90261-L). <http://www.sciencedirect.com/science/article/pii/030442039390261L>.
- Mouchet, A.A., François, L.M.L.M., 1996. Sensitivity of a global oceanic carbon cycle model to the circulation and to the fate of organic matter: preliminary results. *Phys. Chem. Earth* 21, 511–516. [http://dx.doi.org/10.1016/S0079-1946\(97\)81150-0](http://dx.doi.org/10.1016/S0079-1946(97)81150-0). <http://www.sciencedirect.com/science/article/pii/S0079194697811500>.
- Müller, S.A.S.A., Joos, F.F., Edwards, N.R.N.R., Stocker, T.F.T.F., 2006. Water mass distribution and ventilation time scales in a cost-efficient, three-dimensional ocean model. *J. Climate* 19, 5479–5499. <http://dx.doi.org/10.1175/JCLI3911.1>. <http://journals.ametsoc.org/doi/abs/10.1175/JCLI3911.1>.
- Müller, S.A.S.A., Joos, F.F., Plattner, G.-K.G.-K., Edwards, N.R.N.R., Stocker, T.F.T.F., 2008. Modeled natural and excess radiocarbon: sensitivities to the gas exchange formulation and ocean transport strength. *Global Biogeochem. Cycles* 22, GB3011. <http://dx.doi.org/10.1029/2007GB003065>. <http://onlinelibrary.wiley.com/doi/10.1029/2007GB003065/abstract>.
- Munhoven, G.G., 2007. Glacial-interglacial rain ratio changes: implications for atmospheric and ocean-sediment interaction. *Deep-Sea Res. II Top. Stud. Oceanogr.* 54, 722–746. <http://dx.doi.org/10.1016/j.dsr2.2007.01.008>. <http://www.sciencedirect.com/science/article/pii/S0967064507000446>.
- Munhoven, G.G., François, L.M.L.M., 1994. Glacial-interglacial changes in continental weathering: possible implications for atmospheric CO<sub>2</sub>. In: Zahn, R.R., Pedersen, T.F.T.F., Kaminski, M.A.M.A., Labeyrie, L.L. (Eds.), *Carbon Cycling in the Glacial Ocean: Constraints on the Ocean's Role in Global Change*. NATO ASI Series I: Global Environmental Change. vol. 17. Springer-Verlag, Berlin, pp. 39–58.
- Munhoven, G.G., François, L.M.L.M., 1996. Glacial-interglacial variability of atmospheric CO<sub>2</sub> due to changing continental silicate rock weathering: a model study. *J. Geophys. Res.* 101, 21423–21437. <http://dx.doi.org/10.1029/96JD01842>.
- Najjar, R.G.R.G., Jin, X.X., Louanchi, F.F., Aumont, O.O., Caldeira, K.K., Doney, S.C.S.C., Dutay, J.-C.J.-C., Follows, M.M., Gruber, N.N., Joos, F.F., Lindsay, K.K., Maier-Reimer, E.E., Matear, R.J.R.J., Matsumoto, K.K., Monfray, P.P., Mouchet, A.A., Orr, J.C.J.C., Plattner, G.-K.G.-K., Sarmiento, J.L.J.L., Schlitzer, R.R., Slater, R.D.R.D., Weirig, M.-F.M.-F., Yamanaka, Y.Y., Yool, A.A., 2007. Impact of circulation on export

- production, dissolved organic matter, and dissolved oxygen in the ocean: results from Phase II of the Ocean Carbon-cycle Model Intercomparison Project (OCMIP-2). *Global Biogeochem. Cycles* 21 <http://dx.doi.org/10.1029/2006GB002857>. <http://onlinelibrary.wiley.com/doi/10.1029/2006GB002857/abstract> GB3007.
- Najjar, R.G.R.G., Sarmiento, J.L.J.L., Toggweiler, J.R.J.R., 1992. Downward transport and fate of organic matter in the ocean: simulations with a general circulation model. *Global Biogeochem. Cycles* 6, 45–76. <http://dx.doi.org/10.1029/91GB02718>. <http://onlinelibrary.wiley.com/doi/10.1029/91GB02718/abstract>.
- Nickelsen, L.L., Keller, D.P.D.P., Oschlies, A.A., 2015. A dynamic marine iron cycle module coupled to the University of Victoria Earth System Model: the Kiel Marine Biogeochemical Model 2 for UVic 2.9. *Geosci. Model Dev.* 8, 1357–1381. <http://dx.doi.org/10.5194/gmd-8-1357-2015>. <http://www.geosci-model-dev.net/8/1357/2015/00003>.
- Nihoul, J.C.J.C.J., 1994. Do not use a simple model when a complex one will do. *J. Mar. Syst.* 5, 401–406. [http://dx.doi.org/10.1016/0924-7963\(94\)90002-7](http://dx.doi.org/10.1016/0924-7963(94)90002-7). <http://www.sciencedirect.com/science/article/pii/S0924796394900027>.
- Norris, R.D.R.D., Turner, S.K.S.K., Hull, P.M.P.M., Ridgwell, A.A., 2013. Marine ecosystem responses to cenozoic global change. *Science* 341, 492–498. <http://dx.doi.org/10.1126/science.1240543>. <http://www.sciencemag.org/content/341/6145/492>.
- Oeschger, H.H., Siegenthaler, U.U., Schotterer, U.U., Gugelmann, A.A., 1975. A box diffusion model to study the carbon dioxide exchange in nature. *Tellus* 27, 168–192. <http://dx.doi.org/10.1111/j.2153-3490.1975.tb01671.x>.
- Oreskes, N.N., Shrader-Frechette, K.K., Belitz, K.K., 1994. Verification, validation, and confirmation of numerical models in the earth sciences. *Science* 263, 641–646. <http://www.jstor.org/stable/2883078>.
- Pagani, M.M., Huber, M.M., Liu, Z.Z., Bohaty, S.M.S.M., Henderiks, J.J., Sijp, W.W., Krishnan, S.S., DeConto, R.M.R.M., 2011. The role of carbon dioxide during the onset of Antarctic glaciation. *Science* 334, 1261–1264. <http://dx.doi.org/10.1126/science.1203909>. <http://science.sciencemag.org/content/334/6060/1261>.
- Pälike, H.H., Lyle, M.W.M.W., Nishi, H.H., Raffi, I.I., Ridgwell, A.A., Gamage, K.K., Klaus, A.A., Acton, G.G., Anderson, L.L., Backman, J.J., Baldauf, J.J., Beltran, C.C., Bohaty, S.M.S.M., Bown, P.P., Busch, P.P., Channell, J.E.T.J.E.T., Chun, C.O.J.C.O.J., Delaney, M.M., Dewangan, P.P., Dunkley Jones, T.T., Edgar, K.M.K.M., Evans, H.H., Fitch, P.P., Foster, G.L.G.L., Gussone, N.N., Hasegawa, H.H., Hathorne, E.C.E.C., Hayashi, H.H., Herrle, J.O.J.O., Holbourn, A.A., Hovan, S.S., Hyeong, K.K., Iijima, K.K., Ito, T.T., Kamikuri, S.-i.S.-i., Kimoto, K.K., Kuroda, J.J., Leon-Rodriguez, L.L., Malinverno, A.A., Moore Jr T.C., T.C., Murphy, B.H.B.H., Murphy, D.P.D.P., Nakamura, H.H., Ogane, K.K., Ohneiser, C.C., Richter, C.C., Robinson, R.R., Rohling, E.J.E.J., Romero, O.O., Sawada, K.K., Scher, H.H., Schneider, L.L., Sluigs, A.A., Takata, H.H., Tian, J.J., Tsujimoto, A.A., Wade, B.S.B.S., Westerhold, T.T., Wilkens, R.R., Williams, T.T., Wilson, P.A.P.A., Yamamoto, Y.Y., Yamamoto, S.S., Yamazaki, T.T., Zeebe, R.E.R.E., 2012. A cenozoic record of the equatorial pacific carbonate compensation depth. *Nature* 488, 609–614. <http://dx.doi.org/10.1038/nature11360>. <http://www.nature.com/nature/journal/v488/n7413/abs/nature11360.html>.
- Panchuk, K.K., Ridgwell, A.A., Kump, L.R.L.R., 2008. Sedimentary response to Paleocene-Eocene thermal maximum carbon release: a model-data comparison. *Geology* 36, 315–318. <http://dx.doi.org/10.1130/G24474A.1>. <http://geology.gsapubs.org/content/36/4/315>.
- Parekh, P.P., Joos, F.F., Müller, S.A.S.A., 2008. A modeling assessment of the interplay between aeolian iron fluxes and iron-binding ligands in controlling carbon dioxide fluctuations during Antarctic warm events. *Paleoceanography* 23, PA4202. <http://dx.doi.org/10.1029/2007PA001531>. <http://onlinelibrary.wiley.com/doi/10.1029/2007PA001531/abstract>.
- Passow, U.U., 2004. Switching perspectives: do mineral fluxes determine particulate organic carbon fluxes or vice versa? *Geochem. Geophys. Geosyst.* 5 <http://dx.doi.org/10.1029/2003GC000670>. <http://onlinelibrary.wiley.com/doi/10.1029/2003GC000670/abstract>.
- Pedersen, T.F.T.F., Calvert, S.E.S.E., 1990. Anoxia vs. productivity: what controls the formation of organic-carbon-rich sediments and sedimentary rocks? (1). *AAPG Bulletin* 74, 454–466. <http://archives.datapages.com/data/bulletns/1990-91/data/pg/0074/0004/0000/0454.htm>.
- Penman, D.E.D.E., Hönisch, B.B., Zeebe, R.E.R.E., Thomas, E.E., Zachos, J.C.J.C., 2014. Rapid and sustained surface ocean acidification during the Paleocene-Eocene Thermal Maximum. *Paleoceanography* 29 <http://dx.doi.org/10.1002/2014PA002621>. <http://onlinelibrary.wiley.com/doi/10.1002/2014PA002621/abstract> 2014PA002621, 000032.
- Perry, C.T.C.T., Spencer, T.T., Kench, P.S.P.S., 2008. Carbonate budgets and reef production states: a geomorphic perspective on the ecological phase-shift concept. *Coral Reefs* 27, 853–866. <http://dx.doi.org/10.1007/s00338-008-0418-z>. <http://link.springer.com/article/10.1007/s00338-008-0418-z>.
- Petoukhov, V.V., 1980. A zonal climate model of heat and moisture exchange in the atmosphere over the underlying layers of ocean and land. In: Golitsyn, G.S.G.S., Yaglom, A.M.A.M. (Eds.), *Physics of the Atmosphere and the Problem of Climate*.
- Petoukhov, V.V., Claussen, M.M., Berger, A.A., Crucifix, M.M., Eby, M.M., Eliseev, A.V.A.V., Fichet, T.T., Ganopolski, A.A., Goosse, H.H., Kamenkovich, I.I., Mokhov, I.I.I.I., Montoya, M.M., Mysak, L.A.L.A., Sokolov, A.A., Stone, P.P., Wang, Z.Z., Weaver, A.J.A.J., 2005. EMIC intercomparison project (EMIP-CO2): comparative analysis of EMIC simulations of climate, and of equilibrium and transient responses to atmospheric CO<sub>2</sub> doubling. *Climate Dynam.* 25, 363–385. <http://dx.doi.org/10.1007/s00382-005-0042-3>. <http://link.springer.com/article/10.1007/s00382-005-0042-3>.
- Pianosi, F.F., Wagener, T.T., Rougier, J.J., Freer, J.J., Hall, J.J., 2014. Sensitivity analysis of environmental models: a systematic review with practical workflow. In: *Vulnerability, Uncertainty, and Risk*. American Society of Civil Engineers, pp. 290–299.
- Plattner, G.-K.G.-K., Knutti, R.R., Joos, F.F., Stocker, T.F.T.F., von Bloh, W.W., Brovkin, V.V., Cameron, D.D., Driesschaert, E.E., Dutkiewicz, S.S., Eby, M.M., Edwards, N.R.N.R., Fichet, T.T., Hargreaves, J.C.J.C., Jones, C.D.C.D., Loutre, M.F.M.F., Matthews, H.D.H.D., Mouchet, A.A., Müller, S.A.S.A., Nawrath, S.S., Price, A.A., Sokolov, A.A., Strassmann, K.M.K.M., Weaver, A.J.A.J., 2008. Long-term climate commitments projected with climate-carbon cycle models. *J. Climate* 21, 2721–2751. <http://dx.doi.org/10.1175/2007JCLI1905.1>. <http://journals.ametsoc.org/doi/abs/10.1175/2007JCLI1905.1>.
- Ragueneau, O.O., Tréguer, P.P., Leynaert, A.A., Anderson, R.F.R.F., Brzezinski, M.A.M.A., DeMaster, D.J.D.J., Dugdale, R.C.R.C., Dymond, J.J., Fischer, G.G., François, R.R., Heinze, C.C., Maier-Reimer, E.E., Martin-Jézéquel, V.V., Nelson, D.M.D.M., Quéguiner, B.B., 2000. A review of the Si cycle in the modern ocean: recent progress and missing gaps in the application of biogenic opal as a paleoproductivity proxy. *Global Planet. Change* 26, 317–365. [http://dx.doi.org/10.1016/S0921-8181\(00\)00052-7](http://dx.doi.org/10.1016/S0921-8181(00)00052-7). <http://www.sciencedirect.com/science/article/pii/S0921818100000527>.
- Randall, D.A.D.A., Wood, R.A.R.A., Bony, S.S., Colman, R.R., Fichet, T.T., Fyfe, J.J., Kattsov, V.V., Pitman, A.A., Shukla, J.J., Srinivasan, J.J., et al., 2007. Climate models and their evaluation. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. 323.
- Regaudie-de Gioux, A.A., Duarte, C.M.C.M., 2012. Temperature dependence of planktonic metabolism in the ocean. *Global Biogeochem. Cycles* 26, GB1015. <http://dx.doi.org/10.1029/2010GB003907>. <http://onlinelibrary.wiley.com/doi/10.1029/2010GB003907/abstract>.
- Regnier, P.P., Wollast, R.R., Steefel, C.I.C.I., 1997. Long-term fluxes of reactive species in macrotidal estuaries: estimates from a fully transient, multicomponent reaction-transport model. *Mar. Chem.* 58, 127–145. [http://dx.doi.org/10.1016/S0304-4203\(97\)00030-3](http://dx.doi.org/10.1016/S0304-4203(97)00030-3). <http://www.sciencedirect.com/science/article/pii/S0304420397000303>.
- Ridgwell, A.A., 2005. A Mid Mesozoic Revolution in the regulation of ocean chemistry. *Mar. Geol.* 217, 339–357. <http://dx.doi.org/10.1016/j.margeo.2004.10.036>. <http://www.sciencedirect.com/science/article/pii/S0025322705000575>.
- Ridgwell, A.A., 2007. Interpreting transient carbonate compensation depth changes by marine sediment core modeling. *Paleoceanography* 22 <http://dx.doi.org/10.1029/2006PA001372>. <http://onlinelibrary.wiley.com/doi/10.1029/2006PA001372/abstract> PA4102.
- Ridgwell, A.A., Arndt, S.S., 2014. Why dissolved organic matter: DOC in ancient oceans and past climate change. In: Hansell, D.A.D.A., Carlson, C.A.C.A. (Eds.), *Biogeochemistry of Marine Dissolved Organic Matter*, pp. 1–20.
- Ridgwell, A.A., Hargreaves, J.C.J.C., 2007. Regulation of atmospheric CO<sub>2</sub> by deep-sea sediments in an earth system model. *Global Biogeochem. Cycles* 21 <http://dx.doi.org/10.1029/2006GB002764>. <http://onlinelibrary.wiley.com/doi/10.1029/2006GB002764/abstract> n/a–n/a.
- Ridgwell, A.A., Hargreaves, J.C.J.C., Edwards, N.R.N.R., Annan, J.D.J.D., Lenton, T.M.T.M., Marsh, R.R., Yool, A.A., Watson, A.A., 2007. Marine geochemical data assimilation in an efficient Earth System Model of global biogeochemical cycling. *Biogeosciences* 4, 87–104. <https://hal.archives-ouvertes.fr/hal-00297599> 00090.
- Ridgwell, A.A., Schmidt, D.N.D.N., 2010. Past constraints on the vulnerability of marine calcifiers to massive carbon dioxide release. *Nat. Geosci.* 3, 196–200. <http://dx.doi.org/10.1038/ngeo755>. <http://www.nature.com/ngeo/journal/v3/n3/abs/ngeo755.html>.
- Ridgwell, A.A., Zeebe, R.E.R.E., 2005. The role of the global carbonate cycle in the regulation and evolution of the earth system. *Earth Planet. Sci. Lett.* 234, 299–315. <http://dx.doi.org/10.1016/j.epsl.2005.03.006>. <http://www.sciencedirect.com/science/article/pii/S0012821X05001883>.
- Ridgwell, A.J.A.J., 2001. *Glacial-Interglacial Perturbations in the Global Carbon Cycle*. University of East Anglia. <http://ethos.bl.uk/OrderDetails.do?uin=uk.bl.ethos.365134> Ph.D.
- Ridgwell, A.J.A.J., Kennedy, M.J.M.J., Caldeira, K.K., 2003a. Carbonate deposition, climate stability, and Neoproterozoic ice ages. *Science* 302, 859–862. <http://dx.doi.org/10.1126/science.1088342>. <http://science.sciencemag.org/content/302/5646/859>.
- Ridgwell, A.J.A.J., Watson, A.J.A.J., Archer, D.E.D.E., 2002. Modeling the response of the oceanic Si inventory to perturbation, and consequences for atmospheric CO<sub>2</sub>. *Global Biogeochem. Cycles* 16, 1071. <http://dx.doi.org/10.1029/2002GB001877>. <http://onlinelibrary.wiley.com/doi/10.1029/2002GB001877/abstract> 00050.
- Ridgwell, A.J.A.J., Watson, A.J.A.J., Maslin, M.A.M.A., Kaplan, J.O.J.O., 2003b. Implications of coral reef buildup for the controls on atmospheric CO<sub>2</sub> since the Last Glacial Maximum. *Paleoceanography* 18, 1083. <http://dx.doi.org/10.1029/2003PA000893>. <http://onlinelibrary.wiley.com/doi/10.1029/2003PA000893/abstract> 00071.
- Riebesell, U.U., Körtzinger, A.A., Oschlies, A.A., 2009. Sensitivities of marine carbon fluxes to ocean change. *Proc. Natl. Acad. Sci.* 106, 20602–20609. <http://dx.doi.org/10.1073/pnas.0813291106>. <http://www.pnas.org/content/106/49/20602>.
- Roth, R.R., Ritz, S.P.S.P., Joos, F.F., 2014. Burial-nutrient feedbacks amplify the sensitivity of atmospheric carbon dioxide to changes in organic matter remineralisation. *Earth Syst. Dynam.* 5, 321–343. <http://dx.doi.org/10.5194/esd-5-321-2014>. <http://www.earth-syst-dynam.net/5/321/2014/>.
- Ruardij, P.P., Van Raaphorst, W.W., 1995. Benthic nutrient regeneration in the ERSEM ecosystem model of the North Sea. *Neth. J. Sea Res.* 33, 453–483. [http://dx.doi.org/10.1016/0077-5799\(95\)90057-8](http://dx.doi.org/10.1016/0077-5799(95)90057-8). <http://www.sciencedirect.com/science/article/pii/0077579995900578>.
- Sageman, B.B.B.B., Murphy, A.E.A.E., Werne, J.P.J.P., Ver Straeten, C.A.C.A., Hollander, D.J.D.J., Lyons, T.W.T.W., 2003. A tale of shales: the relative roles of production, decomposition, and dilution in the accumulation of organic-rich strata, Middle-Upper Devonian, Appalachian basin. *Chem. Geol.* 195, 229–273. [http://dx.doi.org/10.1016/S0167-4675\(03\)00057-8](http://dx.doi.org/10.1016/S0167-4675(03)00057-8).

- 1016/S0009-2541(02)00397-2. <http://www.sciencedirect.com/science/article/pii/S0009254102003972>.
- Sarmiento, J.L.J.L., Dunne, J.J., Gnanadesikan, A.A., Key, R.M.R.M., Matsumoto, K.K., Slater, R.R., 2002. A new estimate of the CaCO<sub>3</sub> to organic carbon export ratio. *Global Biogeochem. Cycles* 16, 1107. <http://dx.doi.org/10.1029/2002GB001919>. <http://onlinelibrary.wiley.com/doi/10.1029/2002GB001919/abstract>.
- Sarmiento, J.L.J.L., Gruber, N.N., 2006. *Ocean Biogeochemical Dynamics*. 1015 Princeton University Press Princeton.
- Sarmiento, J.L.J.L., Murnane, R.R., Quere, C.L.C.L., Keeling, R.R., Williams, R.G.R.G., 1995. Air-sea CO<sub>2</sub> transfer and the carbon budget of the North Atlantic. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 348, 211–219. <http://dx.doi.org/10.1098/rstb.1995.0063>. <http://rstb.royalsocietypublishing.org/content/348/1324/211>.
- Sarmiento, J.L.J.L., Slater, R.D.R.D., Fasham, M.J.R.M.J.R., Ducklow, H.W.H.W., Toggweiler, J.R.J.R., Evans, G.T.G.T., 1993. A seasonal three-dimensional ecosystem model of nitrogen cycling in the North Atlantic Euphotic Zone. *Global Biogeochem. Cycles* 7, 417–450. <http://dx.doi.org/10.1029/93GB00375>. <http://onlinelibrary.wiley.com/doi/10.1029/93GB00375/abstract>.
- Sarmiento, J.L.J.L., Toggweiler, J.R.J.R., 1984. A new model for the role of the oceans in determining atmospheric pCO<sub>2</sub>. *Nature* 308, 621–624.
- Schmittner, A.A., 2005. Decline of the marine ecosystem caused by a reduction in the Atlantic overturning circulation. *Nature* 434 (7033), 628–633. <http://dx.doi.org/10.1038/nature03476>.
- Schmittner, A.A., Galbraith, E.D.E.D., Hostetler, S.W.S.W., Pedersen, T.F.T.F., Zhang, R.R., 2007. Large fluctuations of dissolved oxygen in the Indian and Pacific oceans during Dansgaard-Oeschger oscillations caused by variations of North Atlantic Deep Water subduction. *Paleoceanography* 22 (3), PA3207. <http://dx.doi.org/10.1029/2006PA001384>. <http://onlinelibrary.wiley.com/doi/10.1029/2006PA001384/abstract>.
- Schmittner, A.A., Lund, D.C.D.C., 2015. Early deglacial Atlantic overturning decline and its role in atmospheric CO<sub>2</sub> rise inferred from carbon isotopes ( $\delta^{13}\text{C}$ ). *Clim. Past* 11, 135–152. <http://dx.doi.org/10.5194/cp-11-135-2015>. <http://www.clim-past.net/11/135/2015/>.
- Schmittner, A.A., Oeschles, A.A., Giraud, X.X., Eby, M.M., Simmons, H.L.H.L., 2005. A global model of the marine ecosystem for long-term simulations: sensitivity to ocean mixing, buoyancy forcing, particle sinking, and dissolved organic matter cycling. *Global Biogeochem. Cycles* 19 (3). <http://dx.doi.org/10.1029/2004GB002283>. <http://onlinelibrary.wiley.com/doi/10.1029/2004GB002283/abstract> n/a–n/a.
- Schmittner, A.A., Somes, C.J.C.J., 2016. Complementary constraints from carbon (<sup>13</sup>C) and nitrogen (<sup>15</sup>N) isotopes on the glacial ocean's soft-tissue biological pump. *Paleoceanography* 31 (6). <http://dx.doi.org/10.1002/2015PA002905>. <http://onlinelibrary.wiley.com/doi/10.1002/2015PA002905/abstract> 2015PA002905.
- Schneider, B.B., Schmittner, A.A., 2006. Simulating the impact of the Panamanian seaway closure on ocean circulation, marine productivity and nutrient cycling. *Earth Planet. Sci. Lett.* 246, 367–380. <http://dx.doi.org/10.1016/j.epsl.2006.04.028>. <http://www.sciencedirect.com/science/article/pii/S0012821X0600330X>.
- Schneider, R.R.R.R., Schulz, H.D.H.D., Hensen, C.C., 2006. Marine carbonates: their formation and destruction. In: Schulz, P.D.H.D.P.D.H.D., Zabel, D.M.D.M. (Eds.), *Marine Geochemistry*. Springer Berlin Heidelberg, pp. 311–337. [http://link.springer.com/chapter/10.1007/3-540-32144-6\\_9](http://link.springer.com/chapter/10.1007/3-540-32144-6_9).
- Sexton, P.F.P.F., Norris, R.D.R.D., Wilson, P.A.P.A., Pälike, H.H., Westerhold, T.T., Röhl, U.U., Bolton, C.T.C.T., Gibbs, S.S., 2011. Eocene global warming events driven by ventilation of oceanic dissolved organic carbon. *Nature* 471, 349–352. <http://dx.doi.org/10.1038/nature09826>. <http://www.nature.com/nature/journal/v471/n7338/abs/nature09826.html>.
- Shaffer, G.G., 1993. Effects of the marine biota on global carbon cycling. In: Heimann, M.M. (Ed.), *The Global Carbon Cycle*. NATO ASI Series Springer Berlin Heidelberg, pp. 431–455. [http://link.springer.com/chapter/10.1007/978-3-642-84608-3\\_18](http://link.springer.com/chapter/10.1007/978-3-642-84608-3_18).
- Shaffer, G.G., 1996. Biogeochemical cycling in the global ocean: 2. New production, redfield ratios, and remineralization in the organic pump. *Journal of Geophysical Research: Oceans* 101, 3723–3745. <http://dx.doi.org/10.1029/95JC03373>. <http://onlinelibrary.wiley.com/doi/10.1029/95JC03373/abstract>.
- Shaffer, G.G., Huber, M.M., Rondanelli, R.R., Pepke Pedersen, J.O.J.O., 2016. Deep time evidence for climate sensitivity increase with warming. *Geophys. Res. Lett.* 43. <http://dx.doi.org/10.1002/2016GL069243>. <http://onlinelibrary.wiley.com/doi/10.1002/2016GL069243/abstract> 2016GL069243.
- Shaffer, G.G., Malskær Olsen, S.S., Pepke Pedersen, J.O.J.O., 2008. Presentation, calibration and validation of the low-order, DCESS earth system model (version 1). *Geosci. Model Dev.* 1, 17–51. <http://dx.doi.org/10.5194/gmd-1-17-2008>. <http://www.geosci-model-dev.net/1/17/2008/>.
- Siegenthaler, U.U., 1983. Uptake of excess CO<sub>2</sub> by an outcrop-diffusion model of the ocean. *J. Geophys. Res.* 88, 3599–3608. <http://dx.doi.org/10.1029/JC088iC06p03599>.
- Siegenthaler, U.U., Wenk, T.T., 1984. Rapid atmospheric CO<sub>2</sub> variations and ocean circulation. *Nature* 308, 624–626.
- Sigman, D.M.D.M., Boyle, E.A.E.A., 2000. Glacial/interglacial variations in atmospheric carbon dioxide. *Nature* 407, 859–869. <http://dx.doi.org/10.1038/35038000>. <http://www.nature.com/nature/journal/v407/n6806/abs/407859a0.html>.
- Sigman, D.M.D.M., Lehman, S.J.S.J., Oppo, D.W.D.W., 2003. Evaluating mechanisms of nutrient depletion and <sup>13</sup>C enrichment in the intermediate-depth Atlantic during the last ice age. *Paleoceanography* 18, 1072. <http://dx.doi.org/10.1029/2002PA000818>. <http://onlinelibrary.wiley.com/doi/10.1029/2002PA000818/abstract>.
- Sigman, D.M.D.M., McCorkle, D.C.D.C., Martin, W.R.W.R., 1998. The calcite lysocline as a constraint on glacial/interglacial low-latitude production changes. *Global Biogeochem. Cycles* 12, 409–427.
- Simmons, C.T.C.T., Matthews, H.D.H.D., Mysak, L.A.L.A., 2015. Deglacial climate, carbon cycle and ocean chemistry changes in response to a terrestrial carbon release. *Climate Dynam.* 1–13. <http://dx.doi.org/10.1007/s00382-015-2646-6>. <http://link.springer.com/article/10.1007/s00382-015-2646-6>.
- Singarayer, J.S.J.S., Richards, D.A.D.A., Ridgwell, A.A., Valdes, P.J.P.J., Austin, W.E.N.W.E.N., Beck, J.W.J.W., 2008. An oceanic origin for the increase of atmospheric radiocarbon during the younger dryas. *Geophys. Res. Lett.* 35, L14707. <http://dx.doi.org/10.1029/2008GL034074>. <http://onlinelibrary.wiley.com/doi/10.1029/2008GL034074/abstract>.
- Sinninghe Damsté, J.S.J.S., Kok, M.D.M.D., Köster, J.J., Schouten, S.S., 1998. Sulfurized carbohydrates: an important sedimentary sink for organic carbon? *Earth Planet. Sci. Lett.* 164, 7–13. [http://dx.doi.org/10.1016/S0012-821X\(98\)00234-9](http://dx.doi.org/10.1016/S0012-821X(98)00234-9). <http://www.sciencedirect.com/science/article/pii/S0012821X98002349>.
- Six, K.D.K.D., Maier-Reimer, E.E., 1996. Effects of plankton dynamics on seasonal carbon flux in an ocean general circulation model. *Global Biogeochem. Cycles* 10, 559–583. <http://dx.doi.org/10.1029/96GB02561>. <http://onlinelibrary.wiley.com/doi/10.1029/96GB02561/abstract>.
- Sluijs, A.A., Zeebe, R.E.R.E., Bijl, P.K.P.K., Bohaty, S.M.S.M., 2013. A middle Eocene carbon cycle conundrum. *Nat. Geosci.* 6, 429–434. <http://dx.doi.org/10.1038/ngeo1807>. <http://www.nature.com/ngeo/journal/v6/n6/abs/ngeo1807.html>.
- Smith, S.S., 1984. Phosphorus versus nitrogen limitation in the marine environment. *LIMNOLOGY*.
- Soetaert, K.K., Herman, P.M.J.P.M.J., 1995. Nitrogen dynamics in the Westerschelde estuary (SW Netherlands) estimated by means of the ecosystem model MOSES. *Hydrobiologia* 311, 225–246. <http://dx.doi.org/10.1007/BF00008583>. <http://link.springer.com/article/10.1007/BF00008583>.
- Soetaert, K.K., Herman, P.M.J.P.M.J., Middelburg, J.J.J.J., 1996. A model of early diagenetic processes from the shelf to abyssal depths. *Geochim. Cosmochim. Acta* 60, 1019–1040. [http://dx.doi.org/10.1016/0016-7037\(96\)00131-0](http://dx.doi.org/10.1016/0016-7037(96)00131-0). <http://www.sciencedirect.com/science/article/pii/0016703796001310>.
- Soetaert, K.K., Hofmann, A.F.A.F., Middelburg, J.J.J.J., Meysman, F.J.R.F.J.R., Greenwood, J.J., 2007. The effect of biogeochemical processes on pH. *Mar. Chem.* 105, 30–51. <http://dx.doi.org/10.1016/j.marchem.2006.12.012>. <http://www.sciencedirect.com/science/article/pii/S0304420306002192>.
- Soetaert, K.K., Middelburg, J.J.J.J., Herman, P.M.J.P.M.J., Buis, K.K., 2000. On the coupling of benthic and pelagic biogeochemical models. *Earth Sci. Rev.* 51, 173–201. [http://dx.doi.org/10.1016/S0012-8252\(00\)00004-0](http://dx.doi.org/10.1016/S0012-8252(00)00004-0). <http://www.sciencedirect.com/science/article/pii/S0012825200000040>.
- Solomatine, D.D., Wagener, T.T., 2011. *Hydrological modeling*. In: Wilderer, P.P. (Ed.), *Treatise on Water Science*. vol. 2. Oxford: Academic Press, pp. 435–457.
- Suess, E.E., 1980. Particulate organic carbon flux in the oceans-surface productivity and oxygen utilization. *Nature* 288, 260–263. <http://dx.doi.org/10.1038/288260a0>. <http://www.nature.com/nature/journal/v288/n5788/abs/288260a0.html>.
- Sun, X.X., Matsumoto, K.K., 2010. Effects of sea ice on atmospheric pCO<sub>2</sub>: a revised view and implications for glacial and future climates. *J. Geophys. Res. Biogeosci.* 115, G02015. <http://dx.doi.org/10.1029/2009JG001023>. <http://onlinelibrary.wiley.com/doi/10.1029/2009JG001023/abstract>.
- Tagliabue, A.A., Aumont, O.O., DeAth, R.R., Dunne, J.P.J.P., Dutkiewicz, S.S., Galbraith, E.E., Misumi, K.K., Moore, J.K.J.K., Ridgwell, A.A., Sherman, E.E., Stock, C.C., Vichi, M.M., Voelker, C.C., Yool, A.A., 2016. How well do global ocean biogeochemistry models simulate dissolved iron distributions? *Global Biogeochem. Cycles* 30, 149–174. <http://dx.doi.org/10.1002/2015GB005289>.
- Tambutté, S.S., Holcomb, M.M., Ferrier-Pagès, C.C., Reynaud, S.S., Tambutté, E.E., Zoccola, D.D., Allemand, D.D., 2011. Coral biomineralization: from the gene to the environment. *J. Exp. Mar. Biol. Ecol.* 408, 58–78. <http://dx.doi.org/10.1016/j.jembe.2011.07.026>. <http://www.sciencedirect.com/science/article/pii/S0022098111003492>.
- Tréguer, P.J.P.J., Rocha, C.L.D.L.C.L.D.L., 2013. The world ocean silica cycle. *Ann. Rev. Mar. Sci.* 5, 477–501. <http://dx.doi.org/10.1146/annurev-marine-121211-172346>.
- Trull, T.T., Bray, S.S., Buesseler, K.K., Lamborg, C.C., Mangani, S.S., Moy, C.C., Valdes, J.J., 2008. In situ measurement of mesopelagic particle sinking rates and the control of carbon transfer to the ocean interior during the vertical flux in the global ocean (VERTIGO) voyages in the north pacific. *Deep-Sea Res. II Top. Stud. Oceanogr.* 55, 1684–1695.
- Tschumi, T.T., Joos, F.F., Gehlen, M.M., Heinze, C.C., 2011. Deep ocean ventilation, carbon isotopes, marine sedimentation and the deglacial CO<sub>2</sub> rise. *Clim. Past* 7, 771–800. <http://dx.doi.org/10.5194/cp-7-771-2011>. <http://www.clim-past.net/7/771/2011/>.
- Tschumi, T.T., Joos, F.F., Parekh, P.P., 2008. How important are Southern Hemisphere wind changes for low glacial carbon dioxide? A model study. *Paleoceanography* 23, PA4208. <http://dx.doi.org/10.1029/2008PA001592>. <http://onlinelibrary.wiley.com/doi/10.1029/2008PA001592/abstract>.
- Tsunogai, S.S., Noriki, S.S., 1991. Particulate fluxes of carbonate and organic carbon in the ocean. Is the marine biological activity working as a sink of the atmospheric carbon? *Tellus B* 43, 256–266. <http://dx.doi.org/10.1034/j.1600-0889.1991.00018.x>. <http://onlinelibrary.wiley.com/doi/10.1034/j.1600-0889.1991.00018.x/abstract>.
- Tyrrell, T.T., 1999. The relative influences of nitrogen and phosphorus on oceanic primary production. *Nature* 400, 525–531. <http://dx.doi.org/10.1038/22941>. <http://www.nature.com/nature/journal/v400/n6744/abs/400525a0.html>.
- Tziperman, E.E., Halevy, I.I., Johnston, D.T.D.T., Knoll, A.H.A.H., Schrag, D.P.D.P., 2011. Biologically induced initiation of neoproterozoic snowball-earth events. *Proc. Natl. Acad. Sci.* 108, 15091–15096. <http://dx.doi.org/10.1073/pnas.1016361108>. <http://www.pnas.org/content/108/37/15091>.
- Uchikawa, J.T., Zeebe, R.E.R.E., 2010. Examining possible effects of seawater pH decline on foraminiferal stable isotopes during the Paleocene-Eocene Thermal Maximum. *Paleoceanography* 25, PA2216. <http://dx.doi.org/10.1029/2009PA001864>. <http://onlinelibrary.wiley.com/doi/10.1029/2009PA001864/abstract>.
- Ushie, H.H., Matsumoto, K.K., 2012. The role of shelf nutrients on glacial-interglacial

- CO<sub>2</sub>: a negative feedback. *Global Biogeochem. Cycles* 26, GB2039. <http://dx.doi.org/10.1029/2011GB004147>. <http://onlinelibrary.wiley.com/doi/10.1029/2011GB004147/abstract>.
- Vecsei, A.A., 2004. A new estimate of global reefal carbonate production including the fore-reefs. *Global Planet. Change* 43, 1–18. <http://dx.doi.org/10.1016/j.gloplacha.2003.12.002>. <http://www.sciencedirect.com/science/article/pii/S0921818104000293>.
- Volk, T.T., Hoffert, M.I.M.I., 1985. Ocean carbon pumps: analysis of relative strengths and efficiencies in ocean-driven atmospheric CO<sub>2</sub> changes. In: Sundquist, E.T.E.T., Broecker, W.S.W.S. (Eds.), *The Carbon Cycle and Atmospheric CO<sub>2</sub>: Natural Variations Archean to Present*. Geophys. Monogr. Ser., vol. 32. AGU, Washington, DC, pp. 99–110.
- Wagner, T.T., Gupta, H.V.H.V., 2005. Model identification for hydrological forecasting under uncertainty. *Stoch. Env. Res. Risk A* 19, 378–387. <http://dx.doi.org/10.1007/s00477-005-0006-5>. <http://link.springer.com/article/10.1007/s00477-005-0006-5>.
- Wakeham, S.G.S.G., Hedges, J.I.J.I., Lee, C.C., Peterson, M.L.M.L., Hernes, P.J.P.J., 1997. Compositions and transport of lipid biomarkers through the water column and surficial sediments of the equatorial Pacific Ocean. *Deep-Sea Res. II Top. Stud. Oceanogr.* 44, 2131–2162. [http://dx.doi.org/10.1016/S0967-0645\(97\)00035-0](http://dx.doi.org/10.1016/S0967-0645(97)00035-0). <http://www.sciencedirect.com/science/article/pii/S0967064597000350>.
- Walker, J.C.G.J.C.G., Opdyke, B.C.B.C., 1995. Influence of variable rates of neritic carbonate deposition on atmospheric carbon dioxide and pelagic sediments. *Paleoceanography* 10, 415–427. <http://dx.doi.org/10.1029/94PA02963>. <http://onlinelibrary.wiley.com/doi/10.1029/94PA02963/abstract>.
- Washington, W.M.W.M., Parkinson, C.C.L.C.C.L., 2005. *An Introduction to Three-Dimensional Climate Modeling*. University Science Books.
- Weber, T.T., Cram, J.A.J.A., Leung, S.W.S.W., DeVries, T.T., Deutsch, C.C., 2016. Deep ocean nutrients imply large latitudinal variation in particle transfer efficiency. *Proc. Natl. Acad. Sci.* 113, 8606–8611. <http://dx.doi.org/10.1073/pnas.1604414113>. <http://www.pnas.org/content/113/31/8606>.
- Welander, P.P., 1959. On the frequency response of some different models describing the transient exchange of matter between the atmosphere and the sea. *Tellus* 11, 348–354. <http://dx.doi.org/10.1111/j.2153-3490.1959.tb00040.x>.
- Wilson, J.D.J.D., Barker, S.S., Ridgwell, A.A., 2012. Assessment of the spatial variability in particulate organic matter and mineral sinking fluxes in the ocean interior: implications for the ballast hypothesis. *Global Biogeochem. Cycles* 26. <http://dx.doi.org/10.1029/2012GB004398>. <http://onlinelibrary.wiley.com/doi/10.1029/2012GB004398/abstract>.
- Winguth, A.M.E.A.M.E., Thomas, E.E., Winguth, C.C., 2012. Global decline in ocean ventilation, oxygenation, and productivity during the Paleocene-Eocene Thermal Maximum: implications for the benthic extinction. *Geology* 40, 263–266. <http://dx.doi.org/10.1130/G32529.1>.
- Yamanaka, Y.Y., Tajika, E.E., 1996. The role of the vertical fluxes of particulate organic matter and calcite in the oceanic carbon cycle: studies using an ocean biogeochemical general circulation model. *Global Biogeochem. Cycles* 10, 361–382. <http://dx.doi.org/10.1029/96GB00634>. <http://onlinelibrary.wiley.com/doi/10.1029/96GB00634/abstract>.
- Yamanaka, Y.Y., Tajika, E.E., 1997. Role of dissolved organic matter in the marine biogeochemical cycle: studies using an ocean biogeochemical general circulation model. *Global Biogeochem. Cycles* 11, 599–612. <http://dx.doi.org/10.1029/97GB02301>. <http://onlinelibrary.wiley.com/doi/10.1029/97GB02301/abstract>.
- Zeebe, R.E.R.E., 2012. LOSCAR: Long-term Ocean-atmosphere-Sediment CARbon cycle Reservoir model v2.0.4. *Geosci. Model Dev.* 5, 149–166. <http://dx.doi.org/10.5194/gmd-5-149-2012>.
- Zeebe, R.E.R.E., Ridgwell, A.A., Zachos, J.C.J.C., 2016. Anthropogenic carbon release rate unprecedented during the past 66 million years. *Nat. Geosci.* 9, 325–329. <http://dx.doi.org/10.1038/ngeo2681>. [http://www.nature.com/ngeo/journal/v9/n4/full/ngeo2681.html?WT.ec\\_id=NGEO-201604&spMailingID=51043954&spUserID=NTQ0MjExMzUxNDgS1&spJobID=883978753&spReportId=ODgzOTc4NzUzS0](http://www.nature.com/ngeo/journal/v9/n4/full/ngeo2681.html?WT.ec_id=NGEO-201604&spMailingID=51043954&spUserID=NTQ0MjExMzUxNDgS1&spJobID=883978753&spReportId=ODgzOTc4NzUzS0).
- Zeebe, R.E.R.E., Wolf-Gladrow, D.D., 2001. *CO<sub>2</sub> in Seawater: Equilibrium, Kinetics, Isotopes: Equilibrium, Kinetics, Isotopes*. Elsevier.
- Zeebe, R.E.R.E., Zachos, J.C.J.C., 2007. Reversed deep-sea carbonate ion basin gradient during Paleocene-Eocene thermal maximum. *Paleoceanography* 22, PA3201. <http://dx.doi.org/10.1029/2006PA001395>. <http://onlinelibrary.wiley.com/doi/10.1029/2006PA001395/abstract>.
- Zeebe, R.E.R.E., Zachos, J.C.J.C., 2013. Long-term legacy of massive carbon input to the Earth system: Anthropocene versus Eocene. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 371, 20120006. <http://dx.doi.org/10.1098/rsta.2012.0006>. <http://rsta.royalsocietypublishing.org/content/371/2001/20120006>.
- Zeebe, R.E.R.E., Zachos, J.C.J.C., Dickens, G.R.G.R., 2009. Carbon dioxide forcing alone insufficient to explain Palaeocene-Eocene Thermal Maximum warming. *Nat. Geosci.* 2, 576–580. <http://dx.doi.org/10.1038/ngeo578>. <http://www.nature.com/ngeo/journal/v2/n8/abs/ngeo578.html>.
- Zickfeld, K.K., Eby, M.M., Weaver, A.J.A.J., Alexander, K.K., Cressin, E.E., Edwards, N.R.N.R., Eliseev, A.V.A.V., Feulner, G.G., Fichet, T.T., Forest, C.E.C.E., Friedlingstein, P.P., Goosse, H.H., Holden, P.B.P.B., Joos, F.F., Kawamiya, M.M., Kicklighter, D.D., Kienert, H.H., Matsumoto, K.K., Mikhov, I.I.I.I., Monier, E.E., Olsen, S.M.S.M., Pedersen, J.O.P.J.O.P., Perrette, M.M., Philippon-Berthier, G.G., Ridgwell, A.A., Schlosser, A.A., Schneider Von Deimling, T.T., Shaffer, G.G., Sokolov, A.A., Spahn, R.R., Steinacher, M.M., Tachiiri, K.K., Tokos, K.S.K.S., Yoshimori, M.M., Zeng, N.N., Zhao, F.F., 2013. Long-term climate change commitment and reversibility: an EMIC intercomparison. *J. Climate* 26, 5782–5809. <http://dx.doi.org/10.1175/JCLI-D-12-00584.1>. <http://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-12-00584.1>.